

COMMODITY SPECIFIC FOOD SAFETY GUIDELINES FOR THE PRODUCTION AND HARVEST OF LETTUCE AND LEAFY GREENS

VERSION 10-11 - ARIZONA

Formatted: Font: Bold, Underline, Font color: Custom Color(RGB(46,116,181))

~~AUGUST 18, 2017~~ SEPTEMBER 14, 2018

Formatted: Underline, Font color: Custom Color(RGB(46,116,181))

Authors Note:

This document reflects Commodity Specific Food Safety Guidelines for the Production and Harvest of Leafy Greens for Arizona. It is based on the Commodity Specific Food Safety Guidelines for the Production and Harvest of Leafy Greens accepted for use by the California Leafy Greens Handler Marketing Agreement and contains minor, non-substantive modifications recommended by the Arizona Leafy Greens Marketing Committee. Arizona law supersedes any requirements in this document that may be in conflict.

46	Table of Contents	
47		
48	Glossary	3
49	Acronyms and Abbreviations	9
50	List of Appendices	10
51	Introduction	11
52	Scope	12
53	1. Purpose	15
54	2. Issue: General Requirements	15
55	3. Issue: Records	15
56	4. Issue: Personnel Qualifications and training	16
57	5. Issue: Environmental Assessments	18
58	6. Issue: Water	19
59	7. Issue: Water Usage to Prevent Product Dehydration	20
60	8. Issue: Soil Amendments	28
61	9. Issue: Nonsynthetic Crop Treatments	37
62	10. Issue: Harvest Equipment, Packing Materials, and Buildings	41
63	11. Issue: Harvest Personnel - Direct Contact with Soil and Contaminants during Harvest	44
64	12. Issue: Field and Harvest Personnel - Transfer of Human Pathogens by Workers	44
65	13. Issue: Equipment Facilitated Cross Contamination	46
66	14. Issue: Flooding	47
67	15. Issue: Production Locations - Climatic Conditions and Environment	51
68	16. Issue: Production Locations - Encroachment by Animals and Urban Settings	51
69	17. Issue: Transportation	60
70	18. Detailed Background Guidance Information	60
71		
72		
73		
74		
75		
76		
77		
78		
79		
80		
81		
82		
83		
84		
85		
86		
87		
88		
89		
90		
91		
92		

Accreditation	A rigorous assessment conducted by an independent science-based organization to assure the overall capability and competency of a laboratory and its quality management systems.
Active compost	Compost feedstock that is in the process of being rapidly decomposed and is unstable. Active compost is generating temperatures of at least 50 degrees Celsius (122 degrees Fahrenheit) during decomposition; or is releasing carbon dioxide at a rate of at least 15 milligrams per gram of compost per day, or the equivalent of oxygen uptake.
Adequate / adequately	That which is needed to accomplish the intended purpose in keeping with good public health practice.
Aerosolized	The dispersion or discharge of a substance under pressure that generates a suspension of fine particles in air or other gas.
Agricultural / Compost tea	A water extract of biological materials (such as compost, manure, non-fecal animal by-products, peat moss, pre-consumer vegetative waste, table waste, or yard trimmings), excluding any form of human waste, produced to transfer microbial biomass, fine particulate organic matter, and soluble chemical components into an aqueous phase. Agricultural / Compost teas are held for longer than one hour before application and are considered non-synthetic crop treatments for the purposes of this document.
Agricultural water	Water used in activities covered in these guidelines where water is intended to, or is likely to, contact lettuce/leafy greens or food contact surfaces, including water used in growing activities (including all irrigation water applied using direct water application methods and water used for preparing crop sprays) and in harvesting, packing, and holding activities (including water used for washing or cooling harvested lettuce/leafy greens and water used for preventing dehydration of lettuce/leafy greens).
Animal by-product	Most parts of an animal that do not include muscle meat including organ meat, nervous tissue, cartilage, bone, blood and excrement.
Animal hazard	Feeding, skin, feathers, fecal matter or signs of animal presence in an area to be harvested in sufficient number and quantity to suggest to a reasonable person the crop may be contaminated.
Adenosine Tri-Phosphate (ATP)	A high energy phosphate molecule required to provide energy for cellular function.
Application interval	Means the time between application of an agricultural input (such as a soil amendment) to a growing area and harvest of leafy greens from the growing area where the agricultural input was applied.
ATP test methods	Exploits knowledge of the concentration of ATP as related to viable biomass or metabolic activity; provides an estimate of cleanliness.
Biofertilizers	Fertilizer materials/products that contain microorganisms such

	as bacteria, fungi, and cyanobacteria that shall promote soil biological activities.
Biosolids	Solid, semisolid, or liquid residues generated during primary, secondary, or advanced treatment of domestic sanitary sewage through one or more controlled processes.
Buildings	Any fully- or partially-enclosed building on the farm that is used for storing of food contact surfaces and packaging materials, including minimal structures that have a roof but no walls.
Colony Forming Units (CFU)	Viable micro-organisms (bacteria, yeasts & mold) either consisting of single cells or groups of cells, capable of growth under the prescribed conditions (medium, atmosphere, time and temperature) to develop into visible colonies (colony forming units) which are counted.
Concentrated Animal Feeding Operation (CAFO)	A lot or facility where animals have been, are or will be stabled or confined and fed or maintained for a total of 45 days or more in any 12 month period and crops, vegetation forage growth, or post-harvest residues are not sustained in the normal growing season over any portion of the lot or facility. In addition, there must be more than 1,000 'animal units' (as defined in 40 CFR 122.23) confined at the facility; or more than 300 animal units confined at the facility if either one of the following conditions are met: pollutants are discharged into navigable waters through a man-made ditch, flushing system or other similar man-made device; or pollutants are discharged directly into waters of the United States which originate outside of and pass over, across, or through the facility or otherwise come into direct contact with the animals confined in the operation.
Coliforms	Gram-negative, non-sporeforming, rod-shaped bacteria that ferment lactose to gas. They are frequently used as indicators of process control, but exist broadly in nature.
Co-management	An approach to conserving soil, water, air, wildlife, and other natural resources while simultaneously minimizing microbiological hazards associated with food production.
Composting	Means a process to produce compost in which organic material is decomposed by the actions of microorganisms under thermophilic conditions for a designated period of time (for example, 3 days) at a designated temperature (for example, 131 °F (55 °C)), followed by a curing stage under cooler conditions.
Cross contamination	The transfer of microorganisms, such as bacteria and viruses, from one place to another.
Curing	The final stage of composting, which is conducted after much of the readily metabolized biological material has been decomposed, at cooler temperatures than those in the thermophilic phase of composting, to further reduce pathogens, promote further decomposition of cellulose and lignin, and stabilize composition. Curing may or may not involve insulation, depending on environmental conditions.

Direct water application	Using agricultural water in a manner whereby the water is intended to, or is likely to, contact leafy greens or food contact surfaces during use of the water.
Enterohemorrhagic <i>E. coli</i>	Shiga toxin-producing <i>E. coli</i> clinically associated with bloody diarrhea.
Escherichia coli (E. coli)	<i>Escherichia coli</i> is a common bacteria that lives in the lower intestines of animals (including humans) and is generally not harmful. It is frequently used as an indicator of fecal contamination, but can be found in nature from non-fecal sources.
Fecal coliforms	Coliform bacteria that grow at elevated temperatures and may or may not be of fecal origin. Useful to monitor effectiveness of composting processes. Also called "thermotolerant coliforms."
Flooding	The flowing or overflowing of a field with water outside a producer's control that is reasonably likely to contain microorganisms of significant public health concern and is reasonably likely to cause adulteration of edible portions of fresh produce in that field.
Food contact surface	Those surfaces that contact human food and those surfaces from which drainage, or other transfer, onto the food or onto surfaces that contact the food ordinarily occurs during the normal course of operations. "Food contact surfaces" includes food contact surfaces of equipment and tools used during harvest, packing and holding.
Food safety assessment	A standardized procedure that predicts the likelihood of harm resulting from exposure to chemical, microbial and physical agents in the diet.
Food safety personnel	Person trained in basic food safety principles and/or working under the auspices of a food safety professional.
Food safety professional	Person entrusted with management level responsibility for conducting food safety assessments before food reaches consumers; requires documented training in scientific principles and a solid understanding of the principles of food safety as applied to agricultural production; in addition this individual must have successfully completed food safety training at least equivalent to that received under standardized curriculum recognized as adequate by the Food and Drug Administration See appendix B for more details.
Geometric Mean	Mathematical def.: the n-th root of the product of n numbers, or: Geometric Mean = n-th root of $(X_1)(X_2) \dots (X_n)$, where X_1, X_2 , etc. represent the individual data points, and n is the total number of data points used in the calculation. Practical def.: the average of the logarithmic values of a data set, converted back to a base 10 number.
Green waste	"Green Waste" means any plant material that is separated at the point of generation, contains no greater than 1.0 percent of physical contaminants by weight, and meets the requirements of section 17868.5. Green material includes, but is not limited to, yard

	trimmings ("Yard Trimmings" means any wastes generated from the maintenance or alteration of public, commercial or residential landscapes including, but not limited to, yard clippings, leaves, tree trimmings, prunings, brush, and weeds), untreated wood wastes, natural fiber products, and construction and demolition wood waste. Green material does not include food material, biosolids, mixed solid waste, material processed from commingled collection, wood containing lead-based paint or wood preservative, mixed construction or mixed demolition debris. "Separated At The Point of Generation" includes material separated from the solid waste stream by the generator of that material. It may also include material from a centralized facility as long as that material was kept separate from the waste stream prior to receipt by that facility and the material was not commingled with other materials during handling. ¹
Ground water	The supply of fresh water found beneath the earth's surface, usually in aquifers, which supply wells and springs. Ground water does not include any water that meets the definition of surface water.
Harvesting	Activities that are traditionally performed on farms for the purpose of removing leafy greens from the field and preparing them for use as food; does not include activities that transform a raw agricultural commodity into a processed food. Examples of harvesting include cutting (or otherwise separating) the edible portion of the leafy greens from the crop plant and removing or trimming parts, cooling, field coring, gathering, hulling, removing stems, trimming of outer leaves and washing.
Hazard	Any biological, physical, or chemical agent that has the potential to cause illness or injury in the absence of its control.
Hobby Farm	A small farm, or rural residence with 25 or fewer animals per acre that is operated without expectation of being the primary source of income.
Holding	Storage of leafy greens in warehouses, cold storage, etc. including activities performed incidental to storage (e.g., activities performed for safe or effective leafy green storage) as well as activities performed as a practical necessity for leafy green distribution (such as blending and breaking down pallets), but does not include activities that transform the raw commodity into a processed food.
Hydroponic	The growing of plants in nutrient solutions with or without an inert medium (as soil) to provide mechanical support.
Indicator microorganisms	An organism that when present suggests the possibility of contamination or under processing.
Known or reasonably foreseeable hazard	Known or reasonably foreseeable hazard means a biological, physical, and chemical hazard that is known to be, or has the potential to be, associated with the farm or the food.
Leafy greens	Iceberg lettuce, romaine lettuce, green leaf lettuce, red leaf lettuce, butter lettuce, baby leaf lettuce (i.e., immature lettuce or leafy greens), escarole, endive, spring mix, spinach, cabbage (green, red and savoy), kale, arugula and chard.

Manure	Animal excreta, alone or in combination with litter (such as straw and feathers used for animal bedding) for use as a soil amendment.
Microorganisms	Yeasts, molds, bacteria, viruses, protozoa, and microscopic parasites and includes species having public health significance and those subjecting leafy greens to decomposition or that otherwise may cause leafy greens to be adulterated.
Monitor	To conduct a planned sequence of observations or measurements to assess whether a process, point or procedure is under control and, when required, to produce an accurate record of the observation or measurement.
Monthly	Because irrigation schedules and delivery of water is not always in a growers control "monthly" for purposes of water sampling means within 35 days of the previous sample.
Most Probable Number (MPN)	Estimated values that are statistical in nature; a method for enumeration of microbes in a sample, particularly when present in small numbers.
Nonsynthetic crop treatments	Any crop input that contains animal manure, an animal product, and/or an animal by-product that is reasonably likely to contain human pathogens. Includes agricultural or compost teas for the purposes of these guidelines.
Oxidation Reduction Potential (ORP)	An intrinsic property that indicates the tendency of a chemical species to acquire electrons and so be reduced; the more positive the ORP, the greater the species' affinity for electrons.
Packing	Placing leafy greens into a container other than packaging them and also includes activities performed incidental to packing (e.g., activities performed for the safe or effective packing of leafy greens (such as sorting, culling, grading, and weighing or conveying incidental to packing or repacking)).
Parts Per Million (ppm)	Usually describes the concentration of something in water or soil; one particle of a given substance for every 999,999 other particles.
Pathogen	A disease causing agent such as a virus, parasite, or bacteria.
Pest	Any objectionable animals or insects, including birds, rodents, flies, and larvae.
Pooled water	An accumulation of standing water; not free-flowing.
Process authority	A regulatory body, person, or organization that has specific responsibility and knowledge regarding a particular process or method; these authorities publish standards, metrics, or guidance for these processes and/or methods.
Ready to eat (RTE) food (excerpted from USFDA 2005 Model Food Code)	<p>(1) "Ready-to-eat food" means FOOD that:</p> <p style="padding-left: 20px;">(a) Is in a form that is edible without additional preparation to achieve FOOD safety, as specified under one of the following: 3-401.11(A) or (B), § 3-401.12, or § 3-402.11, or as specified in 3-401.11(C); or</p> <p style="padding-left: 20px;">(d) May receive additional preparation for palatability or aesthetic, epicurean, gastronomic, or culinary purposes.</p> <p>(2) "Ready-to-eat food" includes:</p> <p style="padding-left: 20px;">(b) Raw fruits and vegetables that are washed as specified</p>

	under § 3-302.15; (c) Fruits and vegetables that are cooked for hot holding, as specified under § 3-401.13; (e) Plant FOOD for which further washing, cooking, or other processing is not required for FOOD safety, and from which rinds, peels, husks, or shells, if naturally present are removed;
Risk mitigation	Actions to reduce the severity/impact of a risk
Sanitary facility	Includes both toilet and hand-washing stations.
Sanitize	To adequately treat cleaned surfaces by a process that is effective in destroying vegetative cells of microorganisms of public health significance, and in substantially reducing numbers of other undesirable microorganisms, but without adversely affecting the product or its safety for the consumer.
Shipping unit/equipment	Any cargo area used to transport leafy greens on the farm or from the farm to cooling, packing, or processing facilities.
Soil amendment	Elements added to the soil, such as compost, peat moss, or fertilizer, to improve its capacity to support plant life.
Surface water	All water open to the atmosphere (rivers, lakes, reservoirs, streams, impoundments, seas, estuaries, etc.) and all springs, wells, or other collectors that are directly influenced by surface water.
Synthetic crop treatments (chemical fertilizers)	Any crop inputs that may be refined, and/or chemically synthesized and/or transformed through a chemical process (e.g. gypsum, lime, sulfur, potash, ammonium sulfate etc.).
Transporter	The entity responsible for transporting product from the field; LGMA guidelines apply only to shippers and cover production through harvesting.
Ultraviolet Index (UV index)	A measure of the solar ultraviolet intensity at the Earth's surface; indicates the day's exposure to ultraviolet rays. The UV index is measured around noon for a one-hour period and rated on a scale of 0-15.
Validated process	A process that has been demonstrated to be effective through a statistically-based study, literature, or regulatory guidance.
Visitor	Any person (other than personnel) who enters your field/operations with your permission.
Water distribution system	Distribution systems -- consisting of pipes, pumps, valves, storage tanks, reservoirs, meters, fittings, and other hydraulic appurtenances – canals, ditches and rivers -- to carry water from its primary source to a lettuce and leafy green crop.

96 Acronyms and Abbreviations
97
98 AFOs: Animal feeding operations
99 AOAC: AOAC International (formerly the Association of Official Analytical Chemists)
100 BAM: Bacteriological Analytical Manual
101 CAFOs: Concentrated animal feeding operations
102 CSG2: *Commodity Specific Guidance for Leafy Greens and Lettuce, 2nd Edition*
103 CFU: colony forming units
104 cGMP: current good manufacturing practices
105 COA: Certificate of Analysis
106 DL: Detection Limit
107 EHEC: Enterohemorrhagic *E. coli*
108 FDA: Food and Drug Administration
109 GAPS: good agricultural practices
110 GLPs: good laboratory practices
111 HACCP: hazard analysis critical control point
112 MPN: most probable number
113 NGO: nongovernmental organization
114 NRCS: Natural Resources Conservation Service
115 ORP: Oxidation reduction potential
116 PPM: parts per million
117 RTE: ready-to-eat
118 SSOPs: Sanitation Standard Operating Procedures
119 TMECC: Test Methods for the Examination of Composting and Compost USEPA: United States
120 Environmental Protection Agency
121 UV: ultraviolet
122 WHO: World Health Organization
123
124
125
126
127
128
129
130

131 **LIST OF APPENDICES**
132 Appendix A: Sanitary Survey
133 Appendix B: Technical Basis Document
134 Appendix C: Crop Sampling Protocol
135 Appendix D: Kinetics of Microbial Inactivation for Alternative Food Processing Technologies
136 Appendix E: Environmental Health Standards for Composting Operations (California Code of
137 Regulations)
138 Appendix T: Training Guidance and Resources
139 Appendix Z: AZ LGMA Resource Agency Contacts
140

141 **INTRODUCTION**

142
143 In 1998, the U.S. Food and Drug Administration (FDA) issued its “Guide to Minimize Microbial
144 Food Safety Hazards for Fresh Fruits and Vegetables.” The practices outlined in this and other
145 industry documents are collectively known as Good Agricultural Practices or GAPs. GAPs provide
146 general food safety guidance on critical production steps where food safety might be compromised
147 during the growing, harvesting, transportation, cooling, packing and storage of fresh produce. More
148 specifically, GAP guidance alerts fruit and vegetable producers, shippers, packers and processors to
149 the potential microbiological hazards associated with various aspects of the production chain
150 including: land history, adjacent land use, water quality, worker hygiene, pesticide and fertilizer use,
151 equipment sanitation and product transportation. The vast majority of the lettuce/leafy greens
152 industry has adopted GAPs as part of normal production operations. Indeed the majority of
153 lettuce/leafy greens producers undergo either internal or external third-party GAP audits on a regular
154 basis to monitor and verify adherence to their GAPs programs. These audit results are often shared
155 with customers as verification of the producer’s commitment to food safety and GAPs.

156 In 2011, the Food Safety Modernization Act (FSMA) was signed into law. After several years of
157 gathering stakeholder input, the FDA published the final regulations promulgating FSMA
158 requirements including regulation of farming operations for the first time in U.S. history. The
159 *Standards for the Growing, Harvesting, Packing, and Holding of Produce for Human Consumption*
160 (the Produce Safety Rule) is the rule that addresses GAPs for farming operations.

161
162 While the produce industry has an admirable record of providing the general public with safe,
163 nutritious fruits and vegetables, it remains committed to continuous improvement with regard to food
164 safety. In 2004, the FDA published a food safety action plan that specifically requested produce
165 industry leadership in developing the next generation of food safety guidance for fruit and vegetable
166 production. These new commodity-specific guidelines focus on providing guidance that enhances the
167 safe growing, processing, distribution and handling of commodities from the field to the end user.
168 The 1st Edition of these new voluntary guidelines was published by the industry in April 2006.

169 In response to continued concerns regarding the microbial safety of fresh produce, this edition of these
170 guidelines (which focuses solely on production and harvest practices) was prepared to provide more
171 specific and quantitative measures of identified best practices for leafy greens production and harvest.
172 In meeting their commitment to keeping the guidelines up-to-date with new scientific and technical
173 advancements, the leafy greens industry has treated the food safety guidelines as a dynamic document
174 by providing routine opportunities for industry members and other stakeholders to recommend
175 revisions and additions. In addition, the guidelines have been updated to reflect the Produce Safety
176 Rule requirements and peer-reviewed research funded by the Center for Produce Safety.

177
178 A key focus of this revision was to identify, where possible and practical, metrics and measures that
179 could be used to assist the industry with compliance with the guidelines. In preparing this document,
180 metrics were researched for three primary areas: water quality, soil amendments, and environmental
181 assessments/conditions. A three-tier approach was used to identify these metrics in as rigorous a
182 manner as possible:

183 1. A comprehensive literature review was conducted to determine if there was a scientifically
184 valid basis for establishing a metric for the identified risk factor or best practice.

185 2. If the literature research did not identify scientific studies that could support an appropriate
186 metric, standards or metrics from authoritative or regulatory bodies were used to establish a
187 metric.
188 3. If neither scientific studies nor authoritative bodies had allowed for suitable metrics,
189 consensus among industry representatives and/or other stakeholders was sought to establish
190 metrics.

191 In the last 10 years, the focus of food safety efforts has been on the farm, initial cooling and
192 distribution points, and value-added processing operations. Fruit and vegetable processing operations
193 have developed sophisticated food safety programs largely centered on current Good Manufacturing
194 Practices (cGMPs) and the principles of Hazard Analysis Critical Control Point (HACCP) programs.
195 As we develop a greater understanding of food safety issues relative to the full spectrum of supply and
196 distribution channels for fruits and vegetables, it has become clear that the next generation of food
197 safety guidance needs to encompass the entire supply chain.

198 In addition to this document, several supplemental documents have been prepared to explain the
199 rationale for the metrics and assist the producer with activities in the field. These documents include a
200 Technical Basis Document that describes in detail and with appropriate citations the bases for the
201 changes made in this edition of this document, a Sanitary Survey document that describes the
202 processes for assessing the integrity and remediation of water systems, and an example product testing
203 plan. All of these items can be found as Appendices to this document.

204 SCOPE

205 The scope of this document pertains only to fresh and fresh-cut lettuce and leafy greens products. It
206 does not include products commingled with non-produce ingredients (e.g. salad kits which may
207 contain meat, cheese, and/or dressings). Examples of “lettuce/leafy greens” include iceberg lettuce,
208 romaine lettuce, green leaf lettuce, red leaf lettuce, butter lettuce, baby leaf lettuce (i.e., immature
209 lettuce or leafy greens), escarole, endive, spring mix, cabbage (green, red and savoy), kale, arugula,
210 chard, radicchio and spinach. These crops are typically considered lettuce and leafy greens by FDA
211 but may not be similarly defined by other state or federal regulatory bodies. This document is also
212 limited to offering food safety guidance practices consistent with the Produce Safety Rule’s provisions
213 for crops grown under outdoor field growing practices and may not address food safety issues related
214 to hydroponic and/or soil-less media production techniques for lettuce/leafy greens.

215 Lettuce/leafy greens may be harvested mechanically or by hand and are almost always consumed
216 uncooked or raw. Because lettuce/leafy greens may be hand-harvested and hand-sorted for quality,
217 there are numerous “touch points” early in the supply chain and a similar number of “touch points”
218 later in the supply chain as the products are used in foodservice or retail operations. Each of these
219 “touch points” represents a potential opportunity for cross-contamination. For purposes of this
220 document, a “touch point” is any occasion when the food is handled by a worker or contacts an
221 equipment food contact surface.

222 Lettuce/leafy greens present multiple opportunities to employ food safety risk management practices
223 to enhance the safety of lettuce/leafy greens. In the production and harvest of lettuce and leafy greens
224 as raw agricultural commodities, GAPs are commonly employed in order to produce the safest
225 products possible. In a processing operation, the basic principles of cGMPs, HACCP, sanitation and
226 documented operating procedures are commonly employed in order to produce the safest products

228 possible. Lettuce/leafy greens are highly perishable and it is strongly recommended that they be
229 distributed, stored and displayed under refrigeration.

230
231 Safe production, packing, processing, distribution and handling of lettuce/leafy greens depend upon a
232 myriad of factors and the diligent efforts and food safety commitment of many parties throughout the
233 distribution chain. No single resource document can anticipate every food safety issue or provide
234 answers to all food safety questions. These guidelines focus on minimizing only the microbial food
235 safety hazards by providing suggested actions to reduce, control or eliminate microbial contamination
236 of lettuce/leafy greens in the field to fork distribution supply chain.

237 All companies involved in the lettuce/leafy greens farm to table supply chain shall implement the
238 recommendations contained within these guidelines to provide for the safe production and handling of
239 lettuce/leafy greens products from field to fork. Every effort to provide food safety education to
240 supply chain partners should also be made. Together with the commitment of each party along the
241 supply chain to review and implement these guidelines, the fresh produce industry is doing its part to
242 provide a consistent, safe supply of produce to the market.

243
244 These guidelines are intended only to convey the best practices associated with the industry. The
245 Produce Marketing Association, the United Fresh Produce Association, Western Growers, and all
246 other contributors and reviewers make no claims or warranties about any specific actions contained
247 herein. It is the responsibility of any purveyor of food to maintain strict compliance with all local,
248 state and federal laws, rules and regulations. These guidelines are designed to facilitate inquiries and
249 developing information that must be independently evaluated by all parties with regard to compliance
250 with legal and regulatory requirements. The providers of this document do not certify compliance
251 with these guidelines and do not endorse companies or products based upon their use of these
252 guidelines.

253 Differences between products, production processes, distribution and consumption, and the ever-
254 changing state of knowledge regarding food safety make it impossible for any single document to be
255 comprehensive and absolutely authoritative. Users of these guidelines should be aware that scientific
256 and regulatory authorities are periodically revising information regarding best practices in food
257 handling, as well as information regarding potential food safety management issues. Users of this
258 document must bear in mind that as knowledge regarding food safety changes, measures to address
259 those changes will also change as will the emphasis on particular issues by regulators and the
260 regulations themselves. Neither this document nor the measures food producers and distributors
261 should take to address food safety are set in stone.

262 Due to the close association between production blocks and environmentally sensitive areas in many
263 locations, it is recommended to review Appendix Z when any mitigation strategies that may impact
264 these areas are employed. Producers should implement strategies that not only protect food safety but
265 also support co-management. All parties involved with implementing the practices outlined in this
266 document should be aware that these metrics are not meant to be in conflict with or discourage co-
267 management practices and principles.

268
269 Users are encouraged to utilize the services of their trade associations, the U.S. Food and Drug
270 Administration, the Center for Produce Safety, the U.S. Department of Agriculture, the U.S.
271 Environmental Protection Agency, the Center for Disease Control and Prevention, and state
272 agricultural, environmental, academic, wildlife and natural resources management agencies and/or
273 public health authorities.

274 The Sanitary Survey and Technical Basis Document prepared as Appendices to these guidelines are
275 considered to be additional resources. They are intended to provide clarification, assist with
276 interpretation and provide additional guidance as users develop food safety programs based on these
277 Guidelines. They are not intended for measurement or verification purposes.

278 **Lettuce/Leafy Greens Commodity Specific Guidance**
279 **Production & Harvest Unit Operations**
280

281 **1. PURPOSE**

282 The issues identified in this document are based on the core elements of Good Agricultural
283 Practices. The specific recommendations contained herein are intended for lettuce and leafy
284 greens only. If these specific recommendations are effectively implemented this would
285 constitute the best practices for a GAP program for the production and harvest unit operations
286 of lettuce and leafy greens.
287

288 **2. ISSUE: GENERAL REQUIREMENTS**

289 In addition to the area-specific requirements discussed in latter sections, there are several
290 general requirements that are part of an effective best practices program. These requirements
291 are outlined below.
292

293 **The Best Practices Are:**

- 294 • A written Leafy Greens Compliance Plan which specifically addresses the Best
295 Practices of this document shall be prepared. This plan shall address at least the
296 following areas: water, soil amendments, environmental factors, work practices,
297 and field sanitation.
- 298 • Shippers shall have an up to date producers list with contact and location
299 information on file.
- 300 • The shipper shall comply with the requirements of The Public Health Security
301 and Bioterrorism Preparedness and Response Act of 2002 (farms are exempt
302 from the Act) including those requirements for recordkeeping (traceability) and
303 registration.
- 304 • Each producer and shipper shall designate an individual responsible for their
305 operation's food safety program. Twenty-four hour contact information shall be
306 available for this individual in case of food safety emergencies.

307 **3. ISSUE: RECORDS**

308 The best practices below complement, but do not supersede recordkeeping requirements in
309 FDA regulations.
310

311 **The Best Practices Are:**

- 312 • All records must include (as applicable to the record):
 - 313 ○ The name (or an identifier e.g., a number that can be linked to the farm/ranch
314 name) and location of the farm
 - 315 ○ Actual values and observations obtained during monitoring

316 ○ An adequate description (e.g., commodity name / specific variety / brand
317 name and, ~~when available~~, any lot number or other identifier) of the leafy
318 green product applicable to the record

319 ○ The location of the growing area (e.g., a specific field) applicable to the
320 record

321 ○ The date and time of the activity documented

322 • All records must be:

323 ○ Created at the time an activity is performed or observed

324 ○ Accurate, legible, and indelible

325 ○ Dated and signed / initialed by the person (or a member of the crew / team)
326 performing the activity documented (does not include the supervisor of those
327 performing the activity)

328 • All records and documents of policies, procedures, and activities to fulfill
329 requirements related to the Leafy Greens Compliance Plan shall be maintained on-
330 site, at an off-site location, or accessible electronically and shall be available for
331 inspection by the end of the day the audit is conducted.

332 • Existing records (e.g., records that are kept to comply with other federal, state, or
333 local regulations or for any other reason) do not need to be duplicated if they contain
334 all of the required information and satisfy the requirements herein. Existing records
335 may be supplemented as necessary to include all of the required information and
336 satisfy the requirements of this section. Records must be kept in the original,
337 electronically or as true copies (e.g., photocopies, pictures, scanned copies,
338 microfilm, microfiche, or other accurate reproductions of the original records).

339 • All required historical records must be readily available and accessible during the
340 retention period for inspection and copying by the LGMA auditor upon oral or
341 written request, except that you have 24 hours to obtain records you keep offsite and
342 make them available and accessible to the auditors for inspection and copying.

343 • If you use electronic techniques to keep records, or to keep true copies of records, or
344 if you use reduction techniques such as microfilm to keep true copies of records, you
345 must provide the records in a format in which they are accessible and legible.

346 • Records shall be kept for a minimum of two years following the date of issuance or
347 occurrence.

348 • Records that relate to the general adequacy of the equipment or processes or records
349 that relate to analyses, sampling, or action plans being used by a farm, including the
350 results of scientific studies, tests, and evaluations, must be retained at the farm for at
351 least 2 years after the use of such equipment or processes, or records related to
352 analyses, sampling, or action plans, is discontinued.

353 **4. ISSUE: PERSONNEL QUALIFICATIONS AND TRAINING**

354 Adequate training of on-farm and shipper personnel is a critically important element in a
355 successful food safety program. In order to align with federal requirements under the Food
356 Safety Modernization Act (FSMA) and to ensure that all activities prescribed in this

357 document are effectively and adequately implemented, the following minimum training
358 requirements must be maintained and documented:

359

360 **The Best Practices Are:**

361 • All personnel (including temporary, part time, seasonal, and contracted personnel)
362 who handle lettuce / leafy greens or who have contact with food-contact surfaces, or
363 who are engaged in the supervision thereof, must:

364 ○ Receive adequate training, as appropriate to the person's duties, upon hiring,
365 and periodically thereafter, at least once annually.

366 ○ Have a combination of education, training, and experience necessary to
367 perform the person's assigned duties in a manner that ensures compliance
368 with these best practices.

369 • Training must be:

370 ○ Conducted in a manner easily understood by personnel being trained.
371 ○ Repeated as necessary and appropriate based on observations or information
372 indicating that personnel are not meeting standards outlined in these best
373 practices.

374 • Minimum training requirements must include:

375 ○ For all personnel who handle (contact) lettuce/leafy greens or supervise those
376 who do so must receive training that includes the following:

377 ■ Principles of food hygiene and safety.
378 ■ The importance of health and personal hygiene for all personnel and
379 visitors including recognizing symptoms of a health condition that is
380 reasonably likely to result in contamination of lettuce/leafy greens or
381 food-contact surfaces with microorganisms of public health
382 significance.

383 ■ The standards established in these best practices that are applicable to
384 the employee's job responsibilities.

385 ○ For harvest personnel, the training program must also address the following
386 minimum requirements related to harvesting activities:

387 ■ Recognizing lettuce/leafy greens that must not be harvested, including
388 product that may be contaminated with known or reasonably
389 foreseeable hazards.

390 ■ Inspecting harvest containers, harvest equipment, and packaging
391 materials to ensure that they are functioning properly, clean, and
392 maintained so as not to become a source of contamination of
393 lettuce/leafy greens with known or reasonably foreseeable hazards.

394 ■ Correcting problems with harvest containers, harvest equipment, or
395 packaging materials or reporting such problems to the supervisor (or
396 other responsible party), as appropriate to the person's job
397 responsibilities.

398 • At least one supervisor or responsible party (e.g., the food safety professional) for
399 each producer providing leafy green products must have successfully completed food
400 safety training at least equivalent to that received under standardized curriculum
401 recognized as adequate by the FDA.
402 • Establish and keep records of training that document required training of personnel,
403 including the date of training, topics covered, and the person(s) trained. Records must
404 be reviewed, dated, and signed, within a week after the records are made, by a
405 supervisor or responsible party.

406 **5. ISSUE: ENVIRONMENTAL ASSESSMENTS**

407 This section addresses assessments that shall be completed and documented prior to the first
408 seasonal planting, within one week prior to harvesting and during harvest operations. These
409 environmental assessments are intended to identify any issues related to the produce field,
410 adjacent land uses, and/or animal hazards that may present a risk to the production block or
411 crop (see Table 5).

412

413 **The Best Practices Are:**

414 • Prior to the first seasonal planting and within one week prior to harvest, perform
415 and document an environmental risk assessment of the production field and
416 surrounding area. Focus these assessments on evaluating the production field for
417 possible animal hazards or other sources of human pathogens of concern,
418 assessing adjacent land uses for possible sources that might contaminate the
419 production field, and evaluating nearby water sources for the potential of past or
420 present flooding.

421 ○ Assessment of Produce Field

422 ■ Evaluate all produce fields for evidence of animal hazards and/or
423 feces. If any evidence is found, follow procedures identified in
424 the “Production Locations - Encroachment by Animals and Urban
425 Settings.”

426 ■ **Evaluate potential environmental sources of contaminants
427 near production locations after a change in weather conditions
428 or weather events that could impact the original risk
429 assessment of the field or block and follow procedures
430 identified in the “Production Locations - Climatic Conditions
431 and Environment” section below.**

432 ○ Assessment of Adjacent Land Use

433 ■ Evaluate all land and waterways adjacent to all production fields
434 for possible sources of human pathogen of concern. These
435 sources include, but are not limited to, manure storage, compost
436 storage, CAFO’s, grazing/open range areas, surface water,
437 sanitary facilities, and composting operations (see Table 6 for
438 further detail). If any possible uses that might result in produce
439 contamination are present consult with the metrics and refer to
440 Appendix Z.

441 ○ **Assessment of CAFOs**

Formatted: Font: Times New Roman, 11.5 pt

Formatted: List Paragraph, Space After: 8 pt, Don't add
space between paragraphs of the same style, Line spacing:
Multiple 1.08 li

Formatted: Font: Times New Roman, 11.5 pt, Bold,
Underline, Font color: Dark Blue

Formatted: Normal, Bulleted + Level: 3 + Aligned at: 1.48"
+ Tab after: 1.73" + Indent at: 1.73"

Formatted: Normal, Indent: Left: 1.73"

Formatted: Font: (Default) Times New Roman, 11.5 pt,
Bold, Underline, Font color: Blue

Formatted: Font: (Default) Times New Roman, 11.5 pt

Formatted: Normal, Bulleted + Level: 2 + Aligned at: 0.98"
+ Tab after: 1.23" + Indent at: 1.23"

443 ■ Conduct and document a rigorous pre-season environmental
 444 assessment of any Concentrated Animal Feeding Operation
 445 that may impact your operation. Include, to the degree
 446 possible, communication with the CAFO operator and/or
 447 third party operator to document Best Management Practices
 448 (BMPs) within the facility, examination of the CAFO for
 449 locations and risk associated with composting, storage, sick
 450 pens, dead piles and other internal operations, examination of
 451 traffic routes associated with the CAFO and examine settling
 452 and manure ponds for any signs of leakage. Note if the CAFO
 453 drainage or discharge is a possible source of contamination.
 454 ■ Record the approximate number of animals within the CAFO
 455 and the method used to determine.
 456 ■ Conduct and document a pre-harvest assessment that
 457 confirms no changes in pre-season conditions. Note if any
 458 discharge events that may impact your crop or operations;
 459 changes in weather condition or weather events occurred
 460 during the production period.
 461 ■ Water sources that are proximate to a CAFO may pose
 462 additional risk and should be closely evaluated. Refer to
 463 Appendix A.

- 464 ○ Assessment of Historical Land Use
 - 465 ■ To the degree practical, determine and document the historical
 - 466 land uses for production fields and any potential issues from these
 - 467 uses that might impact food safety (i.e., hazardous waste sites,
 - 468 landfills, etc.).
- 469 ○ Assessment of Flooding
 - 470 ■ Evaluate all produce fields for evidence of flooding. If any
 - 471 evidence is found, follow procedures identified in the “Flooding”
 - 472 section below.

473 **6. ISSUE: WATER**

474 Water used for production and harvest operations may contaminate lettuce and leafy greens if
 475 water containing human pathogens comes in direct contact with the edible portions of
 476 lettuce/leafy greens. Contamination may also occur by means of water-to-soil followed by
 477 soil-to-lettuce/leafy greens contact. Irrigation methods may have varying potential to
 478 introduce human pathogens or promote human pathogen growth on lettuce and leafy greens
 479 (Stine *et al.*, 2005).

480 There are several different approaches and values that can be utilized to ensure that water is
 481 of appropriate quality for its intended use. The metrics applied in this edition of the
 482 Commodity Specific Guidance should be considered a starting point in industry efforts to
 483 continuously improve the quality of water used in production of these commodities.

484 The current metrics are intended to provide standards associated with water uses; however, it
 485 is known that various water sources have different microbial qualities, and each source

443 **Formatted**

454 **Formatted:** Font: Bold, Underline, Font color: Custom
 455 Color(RGB(46,116,181))

456 **Formatted:** Font: Times New Roman, 11.5 pt

461 **Formatted:** List Paragraph, Space After: 8 pt, Don't add
 462 space between paragraphs of the same style, Line spacing:
 463 Multiple 1.08 li

464 **Formatted:** Font: Times New Roman, 11.5 pt, Bold,
 465 Underline, Font color: Blue

488 should be monitored accordingly. Typical microbial values associated with various sources
489 can be found in the Sanitary Survey document (Appendix A). During the sanitary survey that
490 is performed prior to each growing season expected microbial values and historical
491 monitoring data should be used to evaluate the quality of the water source.

492

493 **The Best Practices Are:**

494

495

496

497

498

499

500

- A water system description shall be prepared. This description can use maps, photographs, drawings or other means to communicate the location of permanent fixtures and the flow of the water system (including any water captured for reuse.). Permanent fixtures include wells, gates, reservoirs, valves, returns and other above ground features that make up a complete irrigation system should be documented in such a manner as to enable location in the field. Water sources and the production blocks they may serve should be documented.
- Water systems that convey untreated human or animal waste must be separated from conveyances utilized to deliver irrigation water.
- Use irrigation water and water in harvest operations that is of appropriate microbial quality for its intended use; see Table 1 and Decision Trees (1A, 1B and 1C) for specific numerical criteria. Appendix B provides the basis for these water quality metrics.
- Perform a sanitary survey prior to use of water in agricultural operations and if water quality microbial tests are at levels that exceed the numerical values set forth in Table 1. The sanitary survey is described in Appendix A.
- Test water as close to the point-of-use as practical, and if microbial levels are above specific action levels, take appropriate remedial and corrective actions.
- Retain documentation of all test results and/or Certificates of Analysis available for inspection for a period of at least 2 years.

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

- **Do not use water from any uncharacterized (untested) water source.**
- **Do not store manure or compost near sources of irrigation water (see Table 6)**
- **Where risk assessments suggest a need, surface waters passing within 400 feet of a CAFO with more than 80,000 head, must be treated to meet microbial acceptance criteria for Postharvest Water (Table 1) if used in any overhead irrigation application at the field level within two weeks of harvest.**

521

522

523

524

525

526

527

528

529

Formatted: Font: Times New Roman, 11.5 pt

Formatted: Indent: Left: 0.5", Tab stops: 0.75", List tab + Not at 1.25"

Formatted: Font: Bold, Underline, Font color: Custom Color(RGB(46,116,181))

Formatted: Indent: Left: 0.5", Bulleted + Level: 1 + Aligned at: 1" + Tab after: 1.25" + Indent at: 1.25", Tab stops: 0.75", List tab + Not at 1.25"

Formatted: Font: Bold, Underline, Font color: Blue

Formatted: Font: Bold, Underline, Font color: Custom Color(RGB(46,116,181))

Formatted: Space Before: 0 pt, After: 0 pt, Bulleted + Level: 1 + Aligned at: 0.75" + Indent at: 1"

Other Considerations for water

o **Water should be treated with EPA approved sanitizers in accordance with label specifications, guidelines for use and consideration of environmental impacts.**

o Evaluate irrigation methods (drip irrigation, overhead sprinkler, furrow, etc.) for their potential to introduce, support or promote the growth of human pathogens on lettuce and leafy greens. Consider such factors as the potential

530 for depositing soil on the crop, presence of pooled or standing water that
531 attracts animals, etc.

532 o When waters from various sources are combined, consider the potential for
533 pathogen growth in the water.

534 o For surface water sources, consider the impact of storm events on irrigation
535 practices. Bacterial loads in surface water are generally much higher after a
536 storm than normal, and caution shall be exercised when using these waters for
537 irrigation.

538 o Use procedures for storing irrigation pipes and drip tape that reduce or
539 eliminate potential pest infestations. Develop procedures to provide for
540 microbiologically safe use of irrigation pipes and drip tape if a pest
541 infestation does occur.

542 o Reclaimed water shall be subject to applicable state and federal regulations
543 and standards. Use of this water for agricultural purposes must meet the most
544 stringent standard as defined by the following: state and federal regulation or
545 Table 1 of this document. Water sample results and analysis provided by the
546 water district or provider may be utilized as records of water source testing
547 for verification and validation audits.

548

549 7. **ISSUE: WATER USAGE TO PREVENT PRODUCT DEHYDRATION**

550 Lettuce/leafy greens may be sprayed with small amounts of water during machine harvest or
551 in the field container just after harvest to reduce water loss. Water used in harvest operations
552 may contaminate lettuce and leafy greens if there is direct contact of water containing human
553 pathogens with edible portions of lettuce/leafy greens.

554

555 **The Best Practices Are:**

556 • Due to the timing of application of water that directly contacts edible portions of
557 lettuce/leafy greens, assure the water is of appropriate microbial quality (e.g.,
558 meets U.S. EPA microbial standards for drinking water).

559 • Test the water source periodically to demonstrate it is of appropriate microbial
560 quality for its intended purpose (e.g., meets U.S. EPA or WHO microbial
561 standards for drinking water) or assure that it has appropriate disinfection
562 potential as described in Table 1.

563 TABLE 1. WATER USE


Use	Metric	Rationale /Remedial Actions
PRE-HARVEST Foliar Applications Whereby Edible Portions of the Crop ARE Contacted by Water (e.g. overhead sprinkler irrigation, pesticides/fungicide application, etc.)	Target Organism: generic <i>E. coli</i> . Sampling Procedure: 100 mL sample collected aseptically at the point of use; i.e., one sprinkler head per water source for irrigation, water tap for pesticides, etc. Water utilized in preseason irrigation operations may be tested and utilized. Sampling Frequency: One sample per water source shall be collected and tested prior to use if >60 days since last test of the water source. Additional samples shall be collected no less than 18 hr apart and at least monthly during use from points within the distribution system. Municipal & Well Exemption: For wells and municipal water sources, if generic <i>E. coli</i> are below detection limits for five consecutive samples, the sampling frequency may be decreased to no less than once every 180 days and the requirements for 60 and monthly sampling are waived. Closed systems with records to demonstrate that all samples of generic <i>E. coli</i> are below detection limits for the two preceding seasons may decrease sampling to a single sample per season. This exemption is void if there is a significant source or distribution system change.	For any given water source (municipal, well, reclaimed water, reservoir or other surface water), samples for microbial testing shall be taken at a point as close to the point of use as practical (as determined by the sampler, to ensure the integrity of the sample, using sampling methods as prescribed in Table 1) where the water contacts the crop, so as to test both the water source and the water distribution system. In a closed water system (meaning no connection to the outside) water samples may be collected from any point within the system but are still preferred as close to point of use as practical. No less than one sample per month per distribution system is required under these metrics unless a system has qualified for an exemption. If there are multiple potential point-of-use sampling points in a distribution system, then samples shall be taken from different point-of-use locations each subsequent month (randomize or rotate sample locations). Water for pre-harvest, direct edible portion contact shall meet or exceed microbial standards for recreational water, based on a rolling geometric mean of the five most recent samples. However, a rolling geometric mean of five samples is not necessarily required prior to irrigation or harvest. If less than five samples are collected prior to irrigation, the acceptance criteria depends on the number of samples taken. If only one sample has been taken, it must be below 126 CFU/100 mL. Once two samples are taken, a geometric mean can be calculated and the normal acceptance criteria apply. If the acceptance criteria are exceeded during this time period, additional samples may be collected to reach a 5 sample rolling geometric mean (as long as the water has not been used for irrigation). The <i>rolling</i> geometric mean calculation starts after 5 samples have been collected. If the water source has not been tested in the past 60 days, the first water sample shall be tested prior to use, to avoid using a contaminated water source. After the first sample is shown to be within acceptance criteria, subsequent samples shall be collected no less frequently than monthly at points of use within the distribution system. Ideally, pre-harvest water should not contain generic <i>E. coli</i> , but low levels do not necessarily indicate that the water is unsafe. Investigation and/or remedial action SHOULD be taken when test results are higher than normal, or indicate an upward trend. Investigation and remedial action SHALL be taken when acceptance criteria are exceeded. Remedial Actions: If the rolling geometric mean (n=5) or any one sample exceeds the acceptance criteria, then the water shall not be used whereby edible portions of the crop are contacted by water until remedial actions have been completed and generic <i>E. coli</i> levels are within acceptance criteria: <ul style="list-style-type: none"> Conduct a sanitary survey of water source and distribution system to determine if a contamination source is evident and can be eliminated. Eliminate identified contamination source(s).

	<p>Test Method: FDA BAM method or any U.S. EPA approved or AOAC accredited for quantitative monitoring of water for generic <i>E. coli</i>. Presence/absence testing with a similar limit of detection may be used as well.</p> <p>Acceptance Criteria: ≤ 126 MPN (or CFU*)/100 mL (rolling geometric mean n=5) and ≤ 235 MPN/100mL for any single sample.</p> <p>*for the purposes of water testing, MPN and CFU shall be considered equivalent.</p> <ul style="list-style-type: none"> For wells, perform a sanitary survey and/or treat as described in Appendix A Sanitary Survey. Retest the water after conducting the sanitary survey and/or taking remedial actions to determine if it meets the outlined microbial acceptance criteria for this use. This sample should represent the conditions of the original water system, if feasible this test should be as close as practical to the original sampling point. A more aggressive sampling program (i.e., sampling once per week instead of once per month) shall be instituted if an explanation for the exceedance is not readily apparent. This type of sampling program should also be instituted if an upward trend is noted in normal sampling results. <p>Crop Testing: If water testing indicates that a crop has been directly contacted with water exceeding acceptance criteria, product shall be sampled and tested for <i>E. coli</i> O157:H7 and <i>Salmonella</i> as described in Appendix C, prior to harvest. If crop testing indicates the presence of either pathogen, the crop shall NOT be harvested for human consumption.</p> <p>Records: Information requirements: Each water sample and analysis shall record: the type of water (canal, reservoir, well, etc) date, time and location of the sample and the method of analysis and detection limit. Records of the analysis of source water may be provided by municipalities, irrigation districts or other water providers. All test results and remedial actions shall be documented and available for verification from the grower/shipper who is the responsible party for a period of two years.</p>
PRE-HARVEST Non-foliar Applications Whereby Edible Portions of the Crop are NOT Contacted by Water (e.g., furrow or drip irrigation, dust abatement water; if water is not used in the vicinity of produce, then testing is not necessary)	<p>Target Organism, Sampling Procedure, Sampling Frequency, Test Method and Municipal & Well Exemption: as described for foliar application.</p> <p>Acceptance Criteria: ≤ 126 MPN /100 mL (rolling geometric mean n=5) and ≤ 576 MPN /100 mL for any single sample.</p> <p>Testing and remedial actions for pre-harvest water that does not come in direct contact with edible portions of the crop are the same as for direct contact water, but acceptance criteria are less stringent because of the reduced risk of contact of the edible portion with contamination from water. Acceptance criteria here are derived from U.S. EPA recreational water standards.</p>

HAND WASH and POSTHARVEST WATER -Direct Product Contact or Food Contact Surfaces	<p><u>Microbial Testing</u></p> <p>Target Organism, Sampling Procedure, Test Method and Municipal & Well Exemption: as described for foliar application.</p> <p>Sampling Frequency: One sample per water source shall be collected and tested prior to use if >60 days since last test of the water source. Additional samples shall be collected at intervals of no less than 18 hr and at least monthly during use.</p> <p>Municipal & Well Exemption: For wells and municipal water sources, if generic <i>E. coli</i> are below detection limits for five consecutive samples, the sampling frequency may be decreased to no less than once every 180 days and the requirements for 60 and monthly sampling are waived. Closed systems with records to demonstrate that all samples of generic <i>E. coli</i> are below detection limits for the two preceding seasons may decrease sampling to a single sample per season. This exemption is void if there is a significant source or distribution system change.</p> <p>Acceptance Criteria: Negative or below DL for all samples</p> <p><u>Physical/Chemical Testing</u></p> <p>Target Variable: Water disinfectant (e.g. chlorine or other disinfectant compound, ORP)</p>	<p>Water that directly contacts edible portions of harvested crop, water used for hand washing, or is used on food contact surfaces, such as equipment or utensils, shall meet the Maximum Contaminant Level Goal for <i>E. coli</i> as specified by U.S. EPA or contain an approved disinfectant at sufficient concentration to prevent cross contamination. Microbial or physical/chemical testing shall be performed, as appropriate to the specific operation, to demonstrate that acceptance criteria have been met. No less than one sample per month per distribution system is required under these metrics unless a system has qualified for an exemption.</p> <p>Single Pass vs. Multiple Pass Systems</p> <ul style="list-style-type: none"> • Single pass use – Water must have non-detectable levels of <i>E. coli</i> or breakpoint disinfectant present at point of entry • Multi-pass use – Water must have non-detectable levels of <i>E. coli</i> and/or sufficient disinfectant to ensure returned water has no detectable <i>E. coli</i> (minimally 1 ppm chlorine). <p>Remedial Actions:</p> <p>If any one sample exceeds the acceptance criteria, then the water shall not be used for this purpose until remedial actions have been completed and generic <i>E. coli</i> or disinfectant levels are within acceptance criteria:</p> <ul style="list-style-type: none"> • Conduct a sanitary survey of water source and distribution system to determine if a contamination source is evident and can be eliminated. Eliminate identified contamination source(s) and/or treat with appropriate disinfectants. For wells, perform a sanitary survey and/or treat as described in Appendix A Sanitary Survey. • Retest the water at the same sampling point after conducting the sanitary survey and/or taking remedial actions to determine if it meets the outlined microbial acceptance criteria for this use. <p>For example, if a water sample for water used to clean food contact surfaces has detectable <i>E. coli</i>, STOP using that water system, examine the distribution line and source inlet as described in Appendix A Sanitary Survey, and retest from the same point of use. Continue testing daily for 5 days at the point closest to use, and do not use the water system until it consistently delivers water that is safe, sanitary and of appropriate microbial quality (i.e. Negative result) for the intended use. If any of the five samples taken during the intensive sampling period after corrective actions have been taken have detectable <i>E. coli</i>, repeat remedial actions and DO NOT use that system until the source of contamination can be corrected.</p>
---	---	--


	<p>Multi Pass Water Acceptance Criteria:</p> <ul style="list-style-type: none"> • <u>Chlorine</u> ≥1 ppm free chlorine after application and pH 6.5 – 7.5 OR • ORP ≥ 650 mV, and pH 6.5 – 7.5 • <u>Other approved treatments</u> per product EPA label for human pathogen reduction in water. <p>Testing Procedure:</p> <ul style="list-style-type: none"> • Chemical reaction based colorimetric test, or • Ion specific probe, or • ORP, or • Other as recommended by disinfectant supplier. <p>Testing Frequency: Continuous monitoring (preferred) with periodic verification by titration OR Routine monitoring if the system can be shown to have a low degree of variation.</p>	<p>Records: All test results and remedial actions shall be documented and available for verification from the user of the water for a period of two years.</p>
--	--	---

564 **Figure 1A. Decision Tree for PRE-HARVEST WATER USE – Foliar Applications**
 565 **whereby edible portions of the crop are contacted by water (e.g. overhead irrigation,**
 566 **pesticide/fungicide applications)**



569
570
571
572

Figure 1B. Decision Tree for PRE-HARVEST WATER USE – Non-Foliar Applications
whereby edible portions of the crop are NOT contacted by water (e.g. furrow or drip irrigation, dust abatement water)



573


574
575

Figure 1C. HAND WASH and POSTHARVEST WATER USE – Direct product contact (e.g. re-hydration, core in field, etc.)

For any given water source (municipal, well, reservoir or other surface water):
Water that directly contacts edible portions of harvested crop, water used for handwashing or is used on food contact surfaces shall meet microbial standards set forth in U.S. EPA National Drinking Water Regulations, and/or contain an approved disinfectant at sufficient concentration to prevent cross contamination.

Sampling Frequency: One sample per water source shall be collected and tested prior to use if >60 days since last test of the water source. Additional samples shall be collected no less than 18 hr apart and a least monthly during use. No less than one sample per month per distribution system is required under these metrics unless a system has qualified for an exemption.

- Sample sources as close to the point-of-use as practical using sampling methods as prescribed in Table 1.
- Analyze samples for generic *E. coli* using a MPN methodology. Other EPA-, FDA- or AOAC



576
577
578

579 **8. ISSUE: SOIL AMENDMENTS**

580 Soil amendments are commonly but not always incorporated prior to planting into
581 agricultural soils used for lettuce/leafy greens production to add organic and inorganic
582 nutrients to the soil as well as intended to improve the physical, chemical, or biological
583 characteristics of soil. Human pathogens may persist in animal manures for weeks or even
584 months (Fukushima *et al.* 1999; Gagliardi and Karns 2000). Proper composting of animal
585 manures via thermal treatment will reduce the risk of potential human pathogen survival.
586 However, the persistence of many human pathogens in agricultural soils depends on many
587 factors (soil type, relative humidity, UV index, etc.) and the effects of these factors is under
588 extensive investigation (Jiang *et al.* 2003; Islam *et al.* 2004).

589
590 Field soil contaminated with human pathogens may provide a means of lettuce and leafy
591 greens contamination. Studies of human pathogens conducted in cultivated field vegetable
592 production models point towards a rapid initial die-off from high pathogen populations but a
593 characteristic and prolonged low level survival. Readily detectable survival is typically less
594 than 8 weeks following incorporation, but has been documented to exceed 12 weeks (Jiang *et*
595 *al.* 2001; Islam *et al.* 2005). Recoverable pathogen populations, using highly sensitive
596 techniques, have been reported to persist beyond this period under some test conditions. The
597 detection of introduced pathogens on mature lettuce plants from these low levels of surviving
598 pathogens was not possible, and the risk was concluded to be negligible. Human pathogens
599 do not persist for long periods of time in high UV index and low relative humidity
600 conditions, but may persist for longer periods of time within aged manure or inadequately
601 composted soil amendments. Therefore, establishing suitably conservative pre-plant
602 intervals, appropriate for specific regional and field conditions, is an effective step towards
603 minimizing risk (Suslow *et al.* 2003).
604

605 **The Best Practices Are:**

- 606 • Do not use biosolids as a soil amendment for production of lettuce or leafy
607 greens.
- 608 • DO NOT USE raw manure or soil amendment that contain untreated animal by-
609 products, un-composted, incompletely composted animal manure and/or green
610 waste or non-thermally treated animal manure to fields which will be used for
611 lettuce and leafy green production.
- 612 • See Table 2 and Decision Trees (Figures 2A and 2B) for numerical criteria and
613 guidance for compost and soil amendments used in lettuce and leafy greens
614 production fields. The Technical Basis Document (Appendix B) describes the
615 process used to develop these metrics.
- 616 • Any soil amendment that does not contain animal manure or other animal by-
617 products must have a document (e.g., ingredient list, statement of identity, letter
618 of guaranty, etc.) from the producer or seller demonstrating that it is manure /
619 animal product free. This document must indicate in some way that manure is
620 not an ingredient used in the production of the amendment or provide the
621 ingredients of the product. A statement of identity or product is sufficient for
622 single-chemical amendments (i.e., “calcium carbonate” or “gypsum”). If “inert
623 ingredients” are listed as part of an amendment, then a document from the
624 producer or seller is necessary indicating manure has not been added. The

625 manure / animal by-product free document must be available for verification
626 before harvest begins and it must be saved and available for inspection for 2
627 years. A new document is required every two years unless there is a significant
628 process or ingredient change.

- 629 • Implement management plans (e.g., timing of applications, storage location,
630 source and quality, transport, etc.) that significantly reduce the likelihood that soil
631 amendments being used contain human pathogens.
- 632 • Verify that the time and temperature process used during the composting process
633 reduces, controls, or eliminates the potential for human pathogens being carried
634 in the composted materials, as applicable to regulatory requirements.
- 635 • Maximize the time interval between soil amendment application and time to
636 harvest.
- 637 • Implement practices that control, reduce or eliminate likely contamination of
638 lettuce/leafy green fields in close proximity to on-farm stacking of manure.
- 639 • Use soil amendment application techniques that control, reduce or eliminate
640 likely contamination of surface water and/or edible crops being grown in adjacent
641 fields.
- 642 • Segregate equipment used for soil amendment handling, preparation, distribution,
643 applications or use effective means of equipment sanitation before subsequent use
644 that effectively reduce the potential for cross contamination.
- 645 • Minimize the proximity of wind-dispersed or aerosolized sources of
646 contamination (e.g., water and manure piles) that may potentially contact growing
647 lettuce/leafy greens or adjacent edible crops. Segregate equipment used for soil
648 amendment applications or use effective means of equipment sanitation before
649 subsequent use.
- 650 • Compost suppliers and on-farm composting operations shall have written
651 Standard Operating Procedures to prevent cross-contamination of in-process and
652 finished compost with raw materials through equipment, runoff, or wind,
653 including instructions for handling, conveying and storing in-process or finished
654 compost like it is untreated if it becomes contaminated. Producers shall annually
655 obtain proof that these documents exist.
- 656 • Compost operations supplying compost to leafy greens crops shall maintain
657 temperature monitoring and turning records for at least two years. Producers
658 purchasing compost shall annually obtain proof from their supplier that this
659 documentation exists. This applies to composting operations regulated under
660 Title 14 CCR as well as smaller operations that do not fall under Title 14.
- 661 • Perform microbiological testing of soil amendments prior to application (Table
662 2).
- 663 • Retain documentation of all processes and test results by lot (at the supplier)
664 and/or Certificates of Analysis available for inspection for a period of at least two
665 years.

666 TABLE 2. SOIL AMENDMENTS

Amendment	Metric/Rationale
Raw Manure, untreated animal products/by-products or Not Fully Composted green waste and/or Animal Manure Containing Soil Amendments (see composted manure process definition below)	DO NOT USE OR APPLY soil amendments that contain un-composted, incompletely composted or non-thermally treated (e.g., heated) animal manure or animal product/by-products to fields which will be used for lettuce and leafy greens production. If these materials have been applied to a field, wait one year prior to producing leafy greens.

667

Composted Soil Amendments (containing animal manure or animal products/by-products)	<p>Please see Figure 2A: Decision Tree for Use of Composted Soil Amendments.</p> <p>Composting Process Validation:</p> <p><u>Enclosed or within-vessel composting:</u> Active compost must maintain a minimum of 131°F for 3 days</p> <p><u>Windrow composting:</u> Active compost must maintain aerobic conditions for a minimum of 131°F or higher for 15 days or longer, with a minimum of five turnings during this period followed by adequate curing.</p> <p><u>Aerated static pile composting:</u> Active compost must be covered with at least 12 inches of insulating materials and maintain a minimum of 131°F for 3 days followed by adequate curing.</p> <p>Target Organisms:</p> <ul style="list-style-type: none"> • <i>Fecal coliforms</i> • <i>Salmonella</i> spp • <i>E. coli</i> O157:H7 <p>Acceptance Criteria:</p> <ul style="list-style-type: none"> • <i>Fecal coliforms</i> <1000 MPN/gram • <i>Salmonella</i>: Negative or < DL (<1/ 30 grams) • <i>E. coli</i> O157:H7: Negative or < DL (<1/ 30 grams)
---	---

	<p>Recommended Test Methods:</p> <ul style="list-style-type: none"> • <i>Fecal coliforms</i>: U.S. EPA Method 1680; multiple tube MPN • <i>Salmonella spp</i>: U.S. EPA Method 1682 • <i>E. coli</i> O157:H7: Any laboratory validated method for compost sampling. • Other U.S. EPA, FDA, AOAC, TMECC or, accredited methods may be used as appropriate. <p>Sampling Plan:</p> <ul style="list-style-type: none"> • A composite sample shall be representative and random and obtained as described in the California state regulations.¹ • Sample may be taken by the supplier if trained by a testing laboratory or state authority. • Laboratory must be certified/accredited for microbial testing by a certification or accreditation body.² <p>Testing Frequency:</p> <ul style="list-style-type: none"> • Each lot before application to production fields. A lot is defined as a unit of production equal to or less than 5,000 cubic yards. <p>Application Interval:</p> <ul style="list-style-type: none"> • Must be applied >45 days before harvest <p>Documentation:</p> <ul style="list-style-type: none"> • All test results and/or Certificates of Analysis shall be documented annually and available for verification from the producer (the responsible party) for a period of two years. Records of process control monitoring for on-farm produced soil amendments must be reviewed, dated, and signed, within a week after the records are made by a supervisor or responsible party. <p>Rationale:</p> <ul style="list-style-type: none"> • The microbial metrics and validated processes for compost are based on allowable levels from California state regulations (CCR Title 14 - Chapter 3.1 - Article 7.2007), with the addition of testing for <i>E. coli</i> O157:H7 as microbe of particular concern. The 45-day application interval was deemed appropriate due to the specified multiple hurdle risk reduction approach outlined. Raw manure must be composted with an approved process and pass testing requirements before an application.
--	--

¹ CCR Title 14 - Chapter-Chapter 3.1 – Article 7 – Section 17868.1
<http://www.calrecycle.ca.gov/Laws/Regulations/title14/ch31a5.htm#article7>

² N/A

Soil amendments containing animal manure that has been heat treated or processed by other equivalent methods.

Please see Figure 2B: Decision Tree for Use of Heat Treated Soil Amendments.

Heat Process Validation

- The heat treatment processes applied to the soil amendment containing animal manure shall be done via a process validated to assure that the process is capable of reducing pathogens of human health significance to acceptable levels.

Target Organism:

- *Fecal coliforms*
- *Salmonella* spp
- *E. coli* O157:H7
- *Listeria monocytogenes*

Acceptance Criteria:

- *Fecal coliforms* Negative or < DL per gram
- *Salmonella*: Negative or < DL (<1/ 30 grams)
- *E. coli* O157:H7: Negative or < DL (<1/ 30 grams)
- *Listeria monocytogenes*: Negative or <DL (<1 CFU / 5 grams)

Recommended Test Methods:

- *Fecal coliforms*: U.S. EPA Method 1680; multiple tube MPN
- *Salmonella* spp: U.S. EPA Method 1682
- *E. coli* O157:H7 *Listeria monocytogenes*: Any laboratory validated method for testing soil amendments.
- U.S. EPA, FDA, AOAC, TMECC or, other accredited methods may be used as appropriate

Sampling Plan:

- Extract at least 12 equivolume samples (identify 12 separate locations from which to collect the sub-sample, in case of bagged product 12 individual bags).
- Sample may be taken by the supplier if trained by a testing laboratory or state authority.
- Laboratory must be certified/accredited by annual review of laboratory protocols based on GLPs by a certification or accreditation body.

Testing Frequency:

- Each lot before application to production fields.
 - In lieu of the above analysis requirement a Certificate of Process Validity Issued by a recognized *Process Authority* can be substituted. This certificate will attest to the process

	<p>validity as determined by either a documented (included w/Certificate) inoculated pack study of the standard process or microbial inactivation calculations of organisms of significant risk (included w/Certificate) as outlined in FDA CFSAN publication “Kinetics of Microbial Inactivation for Alternative Food Processing Technologies. Overarching Principles: Kinetics and Pathogens of Concern for All Technologies” (Incorporated for reference in Appendix E Thermal Process Overview)</p> <p>Application Interval:</p> <ul style="list-style-type: none"> • If the heat treatment process used to inactivate human pathogens of significant public health concern that may be found in animal manure containing soil amendments, is validated and meets the microbial acceptance criteria outlined below, then no time interval is needed between application and harvest. • If the heat treatment process used to inactivate human pathogens of significant public health concern that may be found in animal manure containing soil amendments is not validated but will likely significantly reduce microbial populations of human pathogens and meets microbial acceptance criteria outlined above, then a 45 day interval between application and harvest is required. <p>Documentation:</p> <ul style="list-style-type: none"> • All test results and/or Certificates of Analysis and/or Certificates of Process Validation shall be documented and available for verification from the producer who is the responsible party for a period of two years. The suppliers operation should be validated by a process authority and a record maintained by the producer for a period of two years. <p>Rationale:</p> <ul style="list-style-type: none"> • The microbial metrics and validated processes for compost are based on allowable levels from California state regulations (CCR Title 14 - Chapter 3.1 - Article 7 2007), with the addition of testing for <i>E. coli</i> O157:H7 as the microbe of particular concern. A more stringent level of fecal coliform was also included to address the much more controlled nature of soil amendments produced in this manner. The above suggested application interval was deemed appropriate due to the specified multiple hurdle risk reduction approach outlined. Raw manure must be composted with an approved process and pass testing requirements before application. • FDA has established the validity of D-values and Z-values for key pathogens of concern in foods. This method of process validation is currently acceptable to US regulators. Alternatively, results of an inoculated test pack utilizing the specific process is also an acceptable validation of the lethality of the process.
--	--


Soil Amendments Not Containing Animal Manure

- Any soil amendment that DOES NOT contain animal manure must have documentation that it is manure-free.
- The documentation must be available for verification before harvest begins.
- If there is documentation that the amendment does not contain manure or animal products/by-products then no additional testing is required, and there is no application interval necessary
- Any test results and/or documentation shall be available for verification from the producer who is the responsible party for a period of two years.

669
670
671


Figure 2A. Decision Tree for Composted Soil Amendments (SA)

If raw manure has been directly applied to the field in the past, a 1 year waiting period shall be observed before planting any variety of leafy green crops.



672
673
674
675

676 **Figure 2B. Decision Tree for Heat Treated Animal Manure Containing Soil**
 677 **Amendments (SA)**



678
679

680 9. **ISSUE: NONSYNTHETIC CROP TREATMENTS**

681 Nonsynthetic crop treatments are commonly applied post-emergence for pest and disease
682 control, greening, and to provide organic and inorganic nutrients to the plant during the
683 growth cycle. For the purposes of this document, they are defined as any crop input that
684 contains animal manure, an animal product, and/or an animal by-product that is reasonably
685 likely to contain human pathogens. Due to the potential for human pathogen contamination,
686 these treatments should only be used under conditions that minimize the risk for crop
687 contamination.

688
689 **The Best Practices Are:**

- 690 • Do not use crop treatments that contain raw manure or other untreated animal
691 products or by-products for lettuce or leafy green produce.
- 692 • Do not apply untreated agricultural or compost teas containing added nutrients
693 (e.g., molasses, yeast extract, algal powder, etc.) intended to increase microbial
694 biomass directly to lettuce/leafy greens.
- 695 • Water used to make agricultural teas must meet the water quality requirement for
696 post-harvest water use in Table 1. Liquid crop treatments such as agricultural or
697 compost teas may be used in water distribution systems provided all other
698 requirements herein are met.
- 699 • Retain documentation of all test results available for inspection for a period of at
700 least two years.
- 701 • Implement management plans (e.g. timing of applications, storage location,
702 source and quality, transport, etc.) that assure to the greatest degree practicable
703 that the use of crop treatments does not pose a significant pathogen contamination
704 hazard.
- 705 • Verify that the time and temperature process used during crop treatment
706 manufacture reduces, controls, or eliminates the potential for human pathogens
707 being carried in the nonsynthetic crop treatment materials, as applicable to
708 regulatory requirements.
- 709 • Maximize the time interval between the crop treatment application and time to
710 harvest.
- 711 • Implement practices that control, reduce or eliminate likely contamination of
712 lettuce/leafy green fields that may be in close proximity to on-farm storage of
713 crop treatments (see Table 6 for additional metrics).
- 714 • Use crop treatment application techniques that control, reduce or eliminate the
715 likely contamination of surface water and/or edible crops being grown in adjacent
716 fields.
- 717 • Segregate equipment used for crop treatment applications or use effective means
718 of equipment sanitation before subsequent use.
- 719 • See Table 3 and Decision Tree (Figure 3) for numerical criteria and guidance for
720 nonsynthetic crop treatments used in lettuce and leafy greens production fields.
721 The Technical Basis Document (Appendix B) describes the process used to
722 develop these metrics.


TABLE 3. NONSYNTHETIC CROP TREATMENTS

Treatment	Metric/Rationale
<p><i>Any crop input that contains animal manure, an animal product, and/or an animal by-product that is reasonably likely to contain human pathogens.</i></p> <p>Examples include but are not limited to:</p> <ul style="list-style-type: none"> • Agricultural / Compost teas, • Fish emulsions • Fish meal • Blood meal • "Bio-fertilizers" commonly used for pest control, greening, disease control, fertilizing. <p>Suppliers of these products shall disclose on labels, certificates of analysis, or other companion paperwork whether the product contains any animal manure or products.</p>	<p>Non synthetic crop treatments that contain animal products/by-products or animal manure that have not been physically heat treated or processed by other equivalent methods shall NOT be directly applied to the edible portions of lettuce and leafy greens.</p> <p>Please see Figure 3: Decision Tree for Use of Nonsynthetic Crop Treatments.</p> <p>Process Validation</p> <ul style="list-style-type: none"> • The physical, chemical and/or biological treatment process(es) used to render the crop input safe for application to edible crops must be validated. <p>Target Organism:</p> <ul style="list-style-type: none"> • <i>Fecal coliforms</i> • <i>Salmonella</i> spp • <i>E. coli</i> O157:H7 • <i>Listeria monocytogenes</i> <p>Acceptance Criteria (at point of use):</p> <ul style="list-style-type: none"> • <i>Fecal coliforms</i>: Negative or < DL (<1/ 30 grams or mL) • <i>Salmonella</i>: Negative or < DL (<1/ 30 grams or mL) • <i>E. coli</i> O157:H7: Negative or < DL (<1/ 30 grams or mL) • <i>Listeria monocytogenes</i>: Negative or < DL (< 1 CFU / 5 grams or mL) • Other pathogens appropriate for the source material <p>Recommended Test Methods:</p> <ul style="list-style-type: none"> • <i>Fecal coliforms</i>: U.S. EPA Method 1680; multiple tube MPN • <i>Salmonella</i> spp: U.S. EPA Method 1682 • <i>E. coli</i> O157:H7 and <i>Listeria monocytogenes</i>: Any laboratory validated method for the non synthetic material to be tested. • Other U.S. EPA, FDA, AOAC, TMECC or, accredited methods may be used as appropriate <p>Sampling Plan:</p> <ul style="list-style-type: none"> • 12 point sampling plan composite sample (if solid), one sample per batch if liquid (if liquid-based, then water quality acceptance levels as described in Table 1 should be used) • Sample may be taken by the supplier if trained by the testing laboratory

Treatment	Metric/Rationale
	<ul style="list-style-type: none"> Laboratory must be certified/accredited by annual review of laboratory protocols based on GLPs by a certification or accreditation body. <p>Testing Frequency:</p> <ul style="list-style-type: none"> Each lot before application to production fields. <p>Application Interval:</p> <ul style="list-style-type: none"> If the physical, chemical and/or biological treatment process used to render the crop input safe for application to edible crops is validated and meets that microbial acceptance criteria outlined above, no time interval is needed between application and harvest. If the physical, chemical and/or biological treatment process used to render the crop input safe for application to edible crops is not validated yet meets the microbial acceptance criteria outlined above, a 45 day time interval between application and harvest is required. <p>Documentation:</p> <ul style="list-style-type: none"> All test results and/or Certificates of Analysis shall be documented and available from the producer for verification for a period of 2 years. The producer the party responsible party for maintaining the appropriate records. <p>Rationale:</p> <ul style="list-style-type: none"> The microbial metrics and validated processes for compost are based on allowable levels from California state regulations (CCR Title 14 - Chapter 3.1 - Article 5 2007), with the addition of testing for <i>E. coli</i> O157:H7 as the microbe of particular concern. The above suggested application interval was deemed appropriate due to the specified multiple hurdle risk reduction approach outlined. Any non synthetic crop treatment that contains animal manure must use only fully composted manure in addition to a validated process and pass testing requirements before a application to soils or directly to edible portions of lettuce and leafy greens.

725
726
727

Figure 3. Decision Tree for Nonsynthetic Crop Treatments That Contain Animal Products/by-products



728
729
730
731

732 **Note: Mixtures of soil amendment materials**
733 For soil amendments that contain mixtures of materials each component must meet the
734 requirements of its respective class of materials. The usages allowed will conform to that of
735 the most stringent class of materials utilized in the mixture.

736
737 For example; Soil amendments containing animal manure that has been heat treated or
738 processed by other equivalent methods mixed with soil amendments not containing animal
739 manure would require a process certification for the heat treated or processed by other
740 equivalent methods materials and the components from non-animal manure would require
741 documentation attesting to its manure free status. The resulting mixture could then be
742 applied in accordance with the guidelines associated with the heated treated class of materials
743 (most stringent limits).

744 **10. ISSUE: HARVEST EQUIPMENT, PACKAGING MATERIALS, AND BUILDINGS (FIELD
745 SANITATION)**

746 This section addresses harvest and harvest aid equipment and packaging materials used for
747 lettuce/leafy greens as well as any fully or partially enclosed buildings used to store food
748 contact surfaces and packaging materials. Mechanical or machine harvest has become
749 increasingly prevalent and provides opportunity for increased surface contact exposure. This
750 includes field cored lettuce operations that use various harvest equipment and aids.

751
752 **The Best Practices Are:**

- 753 • Use equipment such as pallets, forklifts, tractors, and vehicles that may have
754 contact with leafy greens in a manner that minimizes the potential for product or
755 food contact surface contamination.
- 756 • **Clean and sanitize food contact surfaces on harvest equipment at the end of
757 each daily harvest.**
- 758 • **Based on inspection, if necessary, rinse and sanitize food contact surfaces on
759 harvest equipment prior to beginning daily harvest.**
- 760 • **All water utilized in cleaning and sanitizing of equipment must meet
761 Postharvest water acceptance criteria.**
- 762 • Prepare an SOP for harvest equipment and containers that addresses the
763 following:
 - 764 ○ **Clean and sanitize when moving between commodities and fields.**
 - 765 ○ Sanitation verification
 - 766 ○ Daily inspection, **cleaning and sanitation**
 - 767 ○ Proper cleaning, sanitation and storage of hand harvest equipment (knives,
768 scythes, etc.)
 - 769 ○ Control procedures when equipment is not in use, including policy for
770 removal of equipment from the work area or site and the use of scabbards,
771 sheathes or other storage equipment.
- 772 • Prepare an SOP for handling and storage of product containers that addresses the
773 following:

Formatted: Font: Bold, Underline, Font color: Custom Color(RGB(46,116,181))

Formatted: List Paragraph, Indent: Left: 0.48", Space Before: 0 pt, After: 8 pt, Don't add space between paragraphs of the same style, Line spacing: Multiple 1.08 li

Formatted: Font: Times New Roman, 11.5 pt, Bold, Underline, Font color: Custom Color(RGB(46,116,181))

Formatted: Font: Bold, Underline, Font color: Custom Color(RGB(46,116,181))

Formatted: List Paragraph, Space Before: 0 pt, After: 8 pt, Don't add space between paragraphs of the same style, Line spacing: Multiple 1.08 li

Formatted: Font: Times New Roman, 11.5 pt, Bold, Underline, Font color: Custom Color(RGB(46,116,181))

Formatted: Font: Bold, Underline, Font color: Custom Color(RGB(46,116,181))

- Overnight storage
- Contact with the ground
- Container assembly (RPC, fiber bin, plastic bin, etc)
- Damaged containers
- Use of containers only as intended
- Prepare an SOP for sanitary operation of equipment which addresses.
 - Spills and leaks
 - Inoperative water sprays
 - Exclusion of foreign objects (including glass, plastic, metal and other debris)
 - Establish and implement cleaning and sanitation schedules for containers and equipment that will be used in hydration.
 - Maintain logs documenting cleaning and sanitation, and retain these records for at least two years.
 - Establish and implement procedures for the storage and control of water tanks and equipment used for hydration operations when not in use.
- Establish and implement appropriate measures that reduce and control the potential introduction of human pathogens at the cut surface during and after mechanical harvest operations. Due to the cut surface being more vulnerable to microbial contamination, this best practice is extremely important and all practical means should be taken to reduce the possibility of introduction of contamination at this process step.
- If re-circulated rinse or antioxidant solutions are used on the cut surface, take all practicable precautions to prevent them from becoming a source of contamination.
- Instruments or controls used to measure, regulate, or record temperatures, hydrogen ion concentration (pH), sanitizer efficacy, or other conditions must be:
 - Accurate and precise as necessary and appropriate for their intended use
 - Adequately maintained; and
 - Adequate in number for their designated uses.

Convey, store, and dispose of trash, litter, and waste to:

- Minimize the potential to attract and harbor pests.
- Protect lettuce/leafy greens, food-contact surfaces, production areas, and agricultural water sources and distribution systems from contamination.

- Seams on food-contact surfaces on equipment and tools must be smoothly bonded or maintained to minimize accumulation of dirt, filth, food particles, and organic materials and the opportunity for harborage or growth of microorganisms.
- Establish sanitation and/or cleaning frequency of food contact and non-food contact surfaces of equipment, tools, and containers by developing and implementing Sanitation Standard Operating Procedures (SSOPs) and a sanitation schedule for machine harvest operations.
- Evaluate the use of cleaning verification methods for harvesting equipment (e.g., ATP test methods).
- Document the date and method of cleaning and sanitizing. A supervisor or responsible party must review, date, and sign these records within a week after the records are made.
- Establish and implement equipment and tool storage and control procedures to minimize the potential for contamination and to prevent it from attracting and harboring pests when not in use.
- Establish policies and implement sanitary design principles that facilitate frequent and thorough cleaning of non-food contact surfaces and cleaning and sanitizing of food contact surfaces.
- Develop and implement appropriate cleaning, sanitizing, storage and handling procedures of all equipment and food contact surfaces to reduce and control the potential for microbial cross contamination.
 - Locate equipment, tool and container cleaning and sanitizing operations away from product and other equipment to reduce the potential for cross contamination.
 - If equipment and tool food contact surfaces have contact with produce that is not covered by the Produce Safety Rule, adequately clean and sanitize before using this equipment to harvest lettuce/leafy greens.
- Allow adequate distance for the turning and manipulation of harvest equipment to prevent cross contamination from areas or adjacent land that may pose a risk.
- Use packaging material that are cleanable or designed for single use and unlikely to support the growth or transfer of bacteria.
- If packaging materials are reused, take steps to ensure food contact surfaces are clean or covered with a clean liner.
- Buildings must be suitable in size, construction and design to facilitate building maintenance and sanitary operations to reduce the potential for contamination of food contact surfaces with known or reasonably foreseeable hazards. Buildings must:
 - Provide sufficient space for placement of equipment and storage of packaging materials.
 - Take proper precautions to reduce potential for contamination of food contact surfaces or packaging materials. Reduce the potential for

854 contamination by effective building design including the separations of
855 operations in which contamination is likely to occur by location, time,
856 partition, enclosed systems, or other effective means.
857 o Provide adequate drainage in all areas where water or other liquid waste is
858 discharged on the ground or floor of the building.
859 o Prevent contamination of food-contact surfaces and packaging materials
860 by protecting them from drips or condensate and excluding pests and
861 animals.

862 **11. ISSUE: HARVEST PERSONNEL - DIRECT CONTACT WITH SOIL AND**
863 **CONTAMINANTS DURING HARVEST (FIELD SANITATION)**

864 After manual harvest of lettuce/leafy greens, placing or stacking product on soil before the
865 product is placed into a container may expose the product to human pathogens if the soil is
866 contaminated. Research has demonstrated that microbes, including human pathogens, can
867 readily attach to cut lettuce/leafy green surfaces (Takeuchi *et al.* 2001).

868
869 **The Best Practices Are:**

870 • Evaluate appropriate measures that reduce and control the potential introduction
871 of human pathogens through soil contact at the cut surface after harvest (e.g.
872 frequency of knife sanitation, no placement of cut surfaces of harvested product
873 on the soil, container sanitation, single use container lining, etc.).
874 • Discard and do not pack any lettuce/leafy greens dropped on the ground during
875 harvest.
876 • Do not stack soiled bins on top of each other if the bottom of one bin has had
877 direct contact with soil unless a protective barrier (*i.e.*, liner, cover, *etc.*) is used
878 to separate the containers.
879 • Establish and implement an SOP for handling in-field trash and other debris
880 including transporting it out of the field in a manner that does not pose a
881 contamination risk.

882 **12. ISSUE: FIELD AND HARVEST PERSONNEL - TRANSFER OF HUMAN PATHOGENS**
883 **BY WORKERS (FIELD SANITATION)**

884 It is possible for persons in the field to transfer microorganisms of significant public health
885 concern to produce during pre-harvest and harvest activities. Establish and implement
886 preventive measures to minimize potential contamination of leafy greens especially during
887 harvest activities when each lettuce/leafy greens plant is touched/handled by harvest crews.
888

889 **The Best Practices Are:**

890 • Use appropriate preventive measures outlined in GAPs such as training in appropriate
891 and effective hand washing, glove use and replacement, and mandatory use of
892 sanitary facilities to reduce and control potential contamination.
893 • Establish and implement a written worker hygiene/practices program (*i.e.*, an SOP)
894 that can be used to verify employee compliance with company food safety policy.
895 This program shall establish the following practices for field and harvest employees
896 as well as visitors.

- 897 ○ During growing and harvesting operations, there must be at least one
898 individual designated as responsible for food safety in compliance with these
899 best practices.
- 900 ○ Use, storage, record keeping, and proper labeling of chemicals
- 901 ○ Follow and be trained in proper sanitation and hygiene practices and policies
902 including:
 - 903 ▪ Requirements for workers to wash their hands with soap and running
904 water before beginning or returning to work, before putting on gloves,
905 after using the toilet, as soon as practical after touching animals or
906 any waste of animal origin and at any other time when hands may
907 have become contaminated.
 - 908 ▪ Requirement for workers' clothing to be clean at the start of the day
909 and appropriate for the operation.
 - 910 ▪ If gloves are used in handling or harvesting lettuce/leafy greens,
911 maintain gloves in an intact and sanitary condition and replace them
912 when no longer able to do so.
 - 913 ▪ Avoiding contact with any animals.
 - 914 ▪ Confinement of smoking, eating and drinking of beverages other than
915 water to designated areas.
 - 916 ▪ Prohibitions on spitting, urinating or defecating in the field.
- 917 ○ Make visitors aware of policies and procedures to protect lettuce/leafy greens
918 and food contact surfaces from contamination by people and take all steps
919 reasonably necessary to ensure that visitors comply with such policies and
920 procedures.
- 921 ▪ Develop and implement a written physical hazard prevention program for leafy green
922 products that are intended for further processing. The program must address the
923 following:
 - 924 ○ Employee clothing and jewelry (head and hair restraints, aprons, gloves,
925 visible jewelry, etc.) Removing or covering hand jewelry (if allowed) that
926 cannot be adequately cleaned and sanitized during periods in which leafy
927 greens are manipulated by hand.
 - 928 ○ Removal of all objects from upper pockets.
 - 929 ○ Designated storage for personal items.
- 930 ▪ Establish and implement a worker health practices program (i.e., an SOP) that
931 address the following issues:
 - 932 ○ Workers with diarrhea disease or symptoms of other infectious disease are
933 prohibited from being in the field and handling fresh produce and food
934 contact surfaces.
 - 935 ○ Workers with open cuts or lesions are prohibited from handling fresh produce
936 and food contact surfaces without specific measures to prevent cross
937 contamination.

938 ○ Actions for employee to take in the event of injury or illness (i.e. notifying a
939 supervisor or other responsible party).

940 ○ A policy describing procedures for handling/disposition of produce or food
941 contact surfaces that have come into contact with blood or other body fluids.

942 ■ A field sanitary facility program (i.e., an SOP) shall be implemented, and it should
943 address the following issues: the number, condition, and placement of field sanitation
944 units according to federal, state or local regulations, the accessibility of the units to
945 the work area, facility maintenance, facility supplies [i.e., hand soap, water (use of
946 antiseptic/sanitizer or wipes, as a substitute for soap and water, is not permitted),
947 single-use paper towels, toilet paper, etc.], facility signage, facility cleaning and
948 servicing, and a response plan for major leaks or spills.

949 ○ During harvest, packing, and holding activities, hand-washing facilities must
950 be furnished with microbial potable running water.

951 ○ Sanitary facilities should be placed such that the location minimizes the
952 impact from potential leaks and/or spills while allowing access for cleaning
953 and service.

954 ○ The location and sanitary design of toilets and hand wash facilities should be
955 optimized to facilitate the control, reduction and elimination of human
956 pathogens from employee hands. Evaluate the location of sanitary facilities
957 to maximize accessibility and use, while minimizing the potential for the
958 facility to serve as a source of contamination.

959 ○ Establish and implement the frequency of sanitary facilities
960 maintenance/sanitation and the appropriate disposal of waste

961 ○ Establish and implement equipment and supply storage and control
962 procedures when not in use.

963 ○ Maintain documentation of maintenance and sanitation schedules and any
964 remedial practices for a period of two years.

965 **13. ISSUE: EQUIPMENT FACILITATED CROSS CONTAMINATION (FIELD**
966 **SANITATION)**

967 When farm equipment has had direct contact with raw untreated manure, untreated compost,
968 waters of unknown quality, animals, or other potential human pathogen reservoirs it may be a
969 source of cross contamination. Such equipment should not be used in proximity to or in
970 areas where it may contact edible portions of lettuce and or leafy greens without proper
971 sanitation.

972 **The Best Practices Are:**

974 ● Identify any field operations that may pose a risk for cross-contamination. These
975 include management personnel in the fields, vehicles used to transport workers,
976 as well as many other possibilities.

977 ● Segregate equipment used in high-risk operations or potentially exposed to high
978 levels of contamination.

979 • Use effective means of equipment cleaning and sanitation before subsequent
980 equipment use in lettuce/leafy greens production, if it was previously used in a
981 high-risk operation.

982 • Develop and implement appropriate means of reducing and controlling the
983 possible transfer of human pathogens to soil and water that may directly contact
984 edible lettuce/leafy green tissues through use of equipment.

985 • Maintain appropriate records related to equipment cleaning and possible cross-
986 contamination issues for a period of two years.

987

988 **14. ISSUE: FLOODING**

989 Flooding for purposes of this document is defined as the flowing or overflowing of a field
990 with water outside of a producer's control, that is reasonably likely to contain
991 microorganisms of significant public health concern and is reasonably likely to cause
992 adulteration of the edible portions of fresh produce in that field. Pooled water (e.g., rainfall)
993 that is not reasonably likely to contain microorganisms of significant public health concern
994 and is not reasonably likely to cause adulteration of the edible portion of fresh produce
995 should not be considered flooding.

996

997 If flood waters contain microorganisms of significant public health concern, crops in close
998 proximity to soil such as lettuce/leafy greens may be contaminated if there is direct contact
999 between flood water or contaminated soil and the edible portions of lettuce/leafy greens
1000 (Wachtel *et al.* 2002a;2002b).

1001

1002 In the November 4, 2005 FDA "Letter to California Firms that Grow, Pack, Process, or Ship
1003 Fresh and Fresh-cut Lettuce/leafy greens" the agency stated that it "considers ready to eat
1004 crops (such as lettuce/leafy greens) that have been in contact with flood waters to be
1005 adulterated due to potential exposure to sewage, animal waste, heavy metals, pathogenic
1006 microorganisms, or other contaminants. FDA is not aware of any method of reconditioning
1007 these crops that will provide a reasonable assurance of safety for human food use or
1008 otherwise bring them into compliance with the law. Therefore, FDA recommends that such
1009 crops be excluded from the human food supply and disposed of in a manner that ensures they
1010 do not contaminate unaffected crops during harvesting, storage or distribution.

1011

1012 "Adulterated food may be subject to seizure under the Federal Food, Drug, and Cosmetic
1013 Act, and those responsible for its introduction or delivery for introduction into interstate
1014 commerce may be enjoined from continuing to do so or prosecuted for having done so. Food
1015 produced under unsanitary conditions whereby it may be rendered injurious to health is
1016 adulterated under § 402(a)(4) of the Federal Food, Drug, and Cosmetic Act (21 U.S.C. 342(a)
1017 (4); (US FDA 2004).

1018

1019 Areas that have been flooded can be separated into three groups: 1) product that has come
1020 into contact with flood water, 2) product that is in proximity to a flooded field but has not
1021 been contacted by flood water, and 3) production ground that was partially or completely
1022 flooded in the past before a crop was planted. The considerations for each situation are
1023 described below and presented in Table 4.

1024

1025 **The Best Practices For Product That Has Come Into Contact With Flood Water**
1026 **Are:**

1027 • See Table 4 for numerical criteria for lettuce and leafy greens production fields
1028 that have possibly come into contact with flood waters. The Technical Basis
1029 Document (Appendix B) describes the process used to develop these metrics.
1030 • FDA considers any crop that has come into contact with floodwater to be an
1031 “adulterated” commodity that cannot be sold for human consumption.
1032 • To reduce the potential for cross contamination do not drive harvest equipment
1033 through flooded areas reasonably likely to contain microorganisms of public
1034 health significance (see previous section).

1035

1036
1037**TABLE 4. FLOODING**

When evidence of flooding in a production block occurs.

Practice	Metric/Rationale
Flooding Defined	The flowing or overflowing of a field with water outside a producer's control that is reasonably likely to contain microorganisms of significant public health concern and is reasonably likely to cause adulteration of edible portions of fresh produce in that field. Additional discussion of this definition and implications for production is provided in the text portion of this document.
Allowable Harvest Distance from Flooding	<ul style="list-style-type: none"> • Buffer and do not harvest any product within 30 ft of the flooding. • Required buffer distance may be greater than 30 ft based on risk analysis by food safety professional. • If there is evidence of flooding, the production block must undergo a detailed food safety assessment by appropriately trained food safety personnel (see Glossary) prior to harvest, as defined in the text of this document.
Verification	<ul style="list-style-type: none"> • Documentation must be archived for a period of two years following the flooding event. Documentation may include photographs, sketched maps, or other means of delineating affected portions of production fields.
Time Interval Before Planting Can Commence Following the Receding of Floodwaters	<ul style="list-style-type: none"> • 60 days prior to planting provided that the soil has sufficient time to dry out. • Appropriate soil testing can be used to shorten this period to 30 days prior to planting. This testing must be performed in a manner that accurately represents the production field and indicates soil levels of microorganisms lower than the recommended standards for processed compost. Suitable representative samples should be collected for the entire area suspected to have been exposed to flooding. For additional guidance on appropriate soil sampling techniques, use the <i>Soil Screening Guidance: Technical Background Document</i> (US EPA 1996). Specifically, Part 4 provides guidance for site investigations. Reputable third-party environmental consultants or laboratories provide sampling services consistent with this guidance. • Appropriate mitigation and mitigation strategies are included in the text portion of the document.
Rationale	<ul style="list-style-type: none"> • The basis for the 30 foot distance is the turnaround distance for production equipment to prevent cross-contamination of non-flooded ground or produce.

1038
1039

1040 **The Best Practices for Product in Proximity to a Flooded Area but Not Contacted**
1041 **By Flood Water Are:**

1042 • Prevent cross contamination between flooded and non-flooded areas (e.g.,
1043 cleaning equipment, eliminating contact of any farming or harvesting equipment
1044 or personnel with the flooded area during growth and harvest of non-flooded
1045 areas).

1046 • To facilitate avoiding contaminated/adulterated produce, place markers
1047 identifying both the high-water line of the flooding and an interval 30 feet beyond
1048 this line. If 30 feet is not sufficient to prevent cross contamination while turning
1049 harvesting or other farm equipment in the field, use a greater appropriate interval.
1050 Take photographs of the area for documentation. Do not harvest product within
1051 the 30 foot buffer zone.

1052
1053 **The Best Practices For Formerly Flooded Production Ground Are:**

1054 • Prior to replanting or soil testing, the designated food safety professional for the
1055 producer shall perform a detailed food safety assessment of the production field.
1056 This designated professional will be responsible for assessing the relative merits
1057 of testing versus observing the appropriate time interval for planting, and also
1058 will coordinate any soil testing plan with appropriate third-party consultants
1059 and/or laboratories that have experience in this type of testing.

1060 • Evaluate the source of flood waters (e.g., drainage canal, river, irrigation canal,
1061 etc.) for potential significant upstream contributors of human pathogens at levels
1062 that pose a significant threat to human health.

1063 • Allow soils to dry sufficiently and be reworked prior to planting subsequent crops
1064 on formerly flooded production ground.

1065 • Do not replant formerly flooded production ground for at least 60 days following
1066 the receding of floodwaters. This period or longer and active tillage of the soil
1067 provide additional protection against the survival of pathogenic organisms.

1068 • If flooding has occurred in the past on the property, soil clearance testing may be
1069 conducted prior to planting leafy greens. Soil testing may be used to shorten the
1070 clearance period to 30 days. If performed, testing must indicate soil levels of
1071 microorganisms lower than the standards for processed compost. Suitable
1072 representative samples should be collected for the entire area suspected to have
1073 been exposed to flooding.

1074 • Sample previously flooded soil for the presence of microorganisms of significant
1075 public health concern or appropriate indicator microorganisms. Microbial soil
1076 sampling can provide valuable information regarding relative risks; however,
1077 sampling by itself does not guarantee that crops grown within the formerly
1078 flooded production area will be free of the presence of human pathogens.

1079 • Evaluate the field history and crop selection on formerly flooded production
1080 ground.

1081 • Assess the time interval between the flooding event, crop planting, and crop
1082 harvest. Comparative soil samples may be utilized to assess relative risk if

1083 significant reductions in indicator microorganisms have occurred within this time
1084 interval.

1085 • Prevent cross-contamination by cleaning or sanitizing any equipment that may
1086 have contacted previously flooded soil (also see the section on Equipment
1087 Facilitated Cross Contamination above).

1088 15. **ISSUE: PRODUCTION LOCATIONS - CLIMATIC CONDITIONS AND ENVIRONMENT**

1089 Lettuce/leafy greens are grown in varying regions but generally in moderate weather
1090 conditions. Cool, humid conditions favor human pathogen persistence (Takeuchi and Frank
1091 2000; Takeuchi *et al.* 2000) while drier climates may present other problems such as
1092 requirements for additional water that may increase the potential for introduction of human
1093 pathogens. Heavy rains in certain areas may also cause lettuce/leafy greens to be exposed to
1094 contaminated soil due to rain splashing. It is important to tailor practices and procedures
1095 designed to promote food safety to the unique environment in which each crop may be
1096 produced

1097

1098 **The Best Practices Are:**

1099 • Consider harvest practices such as removing soiled leaves, not harvesting soiled
1100 heads, etc., when excessive soil or mud builds up on lettuce/leafy greens.

1101 • ~~Take care to reduce the potential for windborne soil, including soil from roads adjacent to fields, water, or other media that may be a source of contamination to come into direct contact with the edible portions of lettuce and leafy greens. Do not allow runoff from adjacent properties to come into contact with produce.~~

1102 • ~~Evaluate and implement practices to reduce the potential for the introduction of pathogens into production blocks by wind or runoff. Such practices may include but are not limited to berms, windbreaks, diversions ditches and vegetated filter strips.~~

1103 • ~~When soil has accumulated on plants, remove soil during the harvest or further processing.~~

1104 **Formatted:** Font: Times New Roman, 11.5 pt

1105 **Formatted:** Bulleted + Level: 1 + Aligned at: 0.25" + Tab after: 0.5" + Indent at: 0.5"

1106 **Formatted:** Left, Space Before: 0 pt, After: 0 pt, No bullets or numbering

1107 **Formatted:** Normal, Justified, Space Before: 6 pt, After: 6 pt, Bulleted + Level: 1 + Aligned at: 0.25" + Tab after: 0.5" + Indent at: 0.5"

1108 **Formatted:** Font: (Default) Arial, 10 pt

1109 **Formatted:** Indent: Left: 0"

1110 **Formatted:** Font: Times New Roman, 11.5 pt

1111 **The Best Practices for Environmental Source of Pathogens and Conditions and Environments:**

1112 • ~~Take care to reduce the potential for windborne soil, including soil from roads adjacent to fields, water, or other media that may be a source of contamination to come into direct contact with the edible portions of lettuce and leafy greens. Do not allow runoff from adjacent properties to come into contact with produce~~

1113 • ~~Evaluate and implement practices to reduce the potential for the introduction of pathogens into production blocks by wind or runoff. Such practices may include but are not limited to berms, windbreaks, diversion ditches and vegetated filter strips~~

1114 • ~~Establish an SOP for production locations that have environmental source of pathogens (i.e. CAFO, dairy, hobby farm and manure or livestock compost facility) and the potential for contamination during weather conditions and events.~~

1115 **Formatted:** List Paragraph, Left, Space Before: 0 pt, After: 8 pt, Don't add space between paragraphs of the same style, Line spacing: Multiple 1.08 li

1116 **Formatted:** Font: Times New Roman, 11.5 pt, Bold

1126

1127 **16. ISSUE: PRODUCTION LOCATIONS - ENCROACHMENT BY ANIMALS AND URBAN**
1128 **SETTINGS**

1129 Lettuce/leafy greens are generally grown in rural areas that may have adjacent wetlands,
1130 wildlands, parks and/or other areas where animals may be present. Some animal species are
1131 known to be potential carriers of various human pathogens (Fenlon 1985; Gorski et al. 2011;
1132 jay et al. 2007; keene et al. 1997; LeJeune et al. 2008; perz et al. 2001). In addition,
1133 extensive development in certain farming communities has also created situations with urban
1134 encroachment and unintentional access by domestic animals and/or livestock which may also
1135 pose varying degrees of risk. Finally, it is possible that some land uses may be of greater
1136 concern than others when located near production fields. Table 6 provides a list of these uses
1137 and recommended buffer distances.

1138

1139 **The Best Practices Are:**

- 1140 • See Tables 5 and 6 and Decision Tree (Figure 5) for numerical criteria and
1141 guidance applicable to animal encroachment and adjacent land uses. The
1142 Technical Basis Document (Appendix B) describes the process used to develop
1143 these metrics.
- 1144 • During the Environmental Assessments discussed in Section 3, the location of
1145 any adjacent land uses that are likely to present a food safety risk should be
1146 documented. In addition, as specified in Table 6, any deviations from the
1147 recommended buffer distances due to mitigation factors or increased risk should
1148 be documented.
- 1149 • Evaluate and monitor animal activity in and proximate to lettuce/leafy greens
1150 fields and production environments. Conduct and document periodic monitoring,
1151 and pre-season, pre-harvest, and harvest assessments. If animals present a
1152 probable risk (medium/high hazard), make particular efforts to reduce their
1153 access to lettuce and leafy green produce.
- 1154 • Fencing, vegetation removal, and destruction of habitat may result in adverse
1155 impacts to the environment. Potential adverse impacts include loss of habitat to
1156 beneficial insects and pollinators; wildlife loss; increased discharges of sediment
1157 and other pollutants resulting from the loss of vegetative filtering; and increased
1158 air quality impacts if bare soil is exposed to wind. It is recommended that
1159 producers check for local, state, and federal laws and regulations that protect
1160 riparian habitat and wetland areas, restrict removal of vegetation or habitat, or
1161 regulate wildlife deterrence measures, including hazing, harassment, lethal and
1162 non-lethal removal, etc.
- 1163 • Evaluate the risk to subsequent crop production or production acreage that has
1164 experienced recent postharvest grazing with or by domesticated animals that used
1165 field culls as a source of animal feed.
- 1166 • Document any probable risk (medium/high hazard) during production and/or
1167 harvest periods and take appropriate corrective action per Table 5 in LGMA
1168 metrics.

1169 • Locate production blocks to minimize potential access by animals and maximize
1170 distances to possible sources of microbial contamination. For example, consider
1171 the proximity to water (i.e., riparian areas), animal harborage, open range lands,
1172 non-contiguous blocks, urban centers, etc. Periodically monitor these factors and
1173 assess during pre-season and pre-harvest assessments as outlined in Tables 5 and
1174 6. If the designated food safety professional deems that there is the potential for
1175 microbial contamination from adjacent areas, a risk assessment shall be
1176 performed to determine the risk level as well as to evaluate potential strategies to
1177 control or reduce the introduction of human pathogens.

1178 • DO NOT harvest areas of fields where unusually heavy activity by animals has
1179 occurred (see Figure 5 Decision Tree).

1180 • If animal intrusions are common on a particular production field, consider
1181 fencing, barriers, noisemakers, and other practices that may reduce intrusions.

1182 • Train harvest employees to recognize and report evidence (e.g., feces) of animal
1183 activity.

1184 • Pooled water (e.g., a seasonal lake) from rainfall may attract animals and should
1185 be considered as part of any land use evaluation.


1186 • Consider controlling risks associated with encroachment by urban development.
1187 Risks may include, but are not limited to, domestic animal fecal contamination of
1188 production fields and harvest equipment and septic tank leaching.

1189 • After a significant event (such as flooding or an earthquake) that could negatively
1190 impact a sewage or septic system, take appropriate steps to ensure that sewage
1191 and septic systems continue to operate in a manner that does not contaminate
1192 produce, food contact surfaces, areas used for produce handling, water sources, or
1193 water distribution systems.

1194 • Producers are encouraged to contact the relevant agencies (e.g., the Regional
1195 Water Quality Control Board and state and federal fish and wildlife agencies) to
1196 confirm the details of these requirements. In addition, producers may wish to
1197 consult with local NRCS to evaluate the food safety risks associated with
1198 wildlife, livestock, domestic animals and other adjacent land uses and to develop
1199 and document strategies to control or reduce the introduction of human pathogens
1200 for each production block.

1201
1202
1203
1204
1205
1206
1207

1208
1209
1210
1211
1212
1213
1214
1215
1216
1217 Figure 5. PRE-HARVEST and HARVEST Assessment – Animal Hazard/Fecal Matter Decision
1218 Tree



Commented [TL1]: Considerations in Assessing Potential Hazards and Risks Associated with Animal Activity in the Field (domestic, wild)

- Volume and concentration of fecal material in the field and production area
- Frequency of animal sightings and sign (e.g., tracks, scat, rubbing, animal damage to crop)
- Animal species likely to aggregate (e.g., flocks and herds) and produce concentrated areas of fecal material and incidental contact with the crop
- Potential for animals, pests, rodents and birds as a risk source to transport pathogens from a high risk source (e.g., CAFO, garbage dump, sewage treatment facility) to the field
- Species with seasonal migrations that result in increased population density and potential for activity in the field

1220
1221

TABLE 5. ANIMAL HAZARD IN FIELD (WILD OR DOMESTIC)
When evidence of animal intrusion in a production block occurs.

Issue	Metric	Remedial Actions
Evidence of Intrusion	<p><u>Frequency</u></p> <ul style="list-style-type: none">• There shall be a periodic monitoring plan in place for production fields.• There shall be Pre-Season, Pre-Harvest, and Harvest Assessments <p><u>Variables</u></p> <ul style="list-style-type: none">• Physical observation of animals in the field• Downed fences• Animal tracks in production block• Animal feces or urine in production block• Damaged or eaten plants in production block	<ul style="list-style-type: none">• If there is evidence of intrusion by animals, the production block must undergo a detailed food safety assessment by appropriately trained food safety personnel (see Glossary) prior to harvest, as defined in the text of this document.• Animal intrusion events shall be categorized as low or medium/high hazard. An example of a low hazard might be a sign of animal intrusion into the leafy green production area by a single animal or solitary bird with minimal to no fecal deposition.• Corrective actions for “Low hazard” animal intrusion shall be carried out according to company SOP.• Corrective actions for “medium/high hazard” animal intrusion shall be carried out per the accepted LGMA metrics and must include food safety buffers and do not harvest areas.• In developing preventive remedial and corrective actions, consider consulting with wildlife and/or domestic animal experts as appropriate.• If remedial actions, such as appropriate no harvest buffers, cannot be formulated to control or eliminate the identified risk, do not harvest and instead destroy the contaminated crop.• Equipment used to destroy crop must be cleaned and sanitized upon exiting the field.• Formulate effective corrective actions. Prior to taking action that may affect natural resources, producers should check local, state and federal laws and regulations that protect riparian habitat and wetland areas, restrict removal of vegetation or habitat, or restrict construction of wildlife deterrent fences in riparian areas or wildlife corridors.• Food safety assessments and corrective actions shall be documented and available for verification for a period of two years.

Issue	Metric	Remedial Actions
Allowable Harvest Distance from Evidence of Intrusion	<p>Please see Figure 5. Decision Tree for Conducting Pre-Harvest and Harvest Assessments.</p> <p><u>Monitoring</u></p>	
	<p>Conduct periodic monitoring and, pre-season, pre-harvest and harvest assessments. Evaluate and monitor animal activity in and proximate to lettuce/leafy greens fields and production environments.</p>	
	<p><u>Pre-Harvest Assessment and Daily Harvest Assessment</u></p> <ul style="list-style-type: none"> Conduct the pre-harvest assessment not more than one week prior to harvest. Conduct the daily harvest assessment on each day of harvest. 	
	<p>Fecal Material</p> <ul style="list-style-type: none"> Do not harvest any produce that has come into direct contact with fecal material. If evidence of fecal material is found, conduct a food safety assessment using qualified personnel. Do not harvest any crop found within a minimum 5 foot radius buffer distance from the spot of the contamination unless remedial action can be found that adequately control the risk. The food safety professional can increase this buffer distance if deemed appropriate. 	
	<p>Intrusion</p> <ul style="list-style-type: none"> If evidence of animal intrusion is found in a production field, conduct a visual food safety assessment to determine whether the intrusion is a probable (medium/high hazard) or negligible (low hazard) risk. Low hazard (negligible risk) can be corrected by following a company SOP. Medium/high hazard (probable risk) intrusion should include a three foot buffer radius where the impacted crop has been isolated. 	
	<p><u>Daily Harvest Assessment ONLY</u></p> <p>If evidence of medium/high hazard risk animal intrusion into the production block is not discovered until harvest operations:</p> <ul style="list-style-type: none"> Stop harvest operations. Initiate an intensified block assessment for evidence of further contamination and take appropriate actions per the aforementioned actions. If evidence of intrusion is discovered during production block harvest operations and the harvest rig has been potentially contaminated by contaminated product or feces, clean and sanitize the equipment before resuming harvest operations. Require all employees to wash and sanitize their hands/gloves before resuming harvest operations. If contamination is discovered in harvest containers such as bins/totes, discard the product, and clean and sanitize the container before reuse. 	
Verification	<ul style="list-style-type: none"> Archive documentation for a period of two years following the intrusion event. Documentation may include photographs, sketched maps, or other means of delineating affected portions of production fields. 	
Rationale	<ul style="list-style-type: none"> The basis of these metrics is qualitative assessment of the relative risk from a variety of intrusions. Some animal feces and some signs of intrusion (feces vs. tracks) are considered to be of more concern than others. Because it is difficult to develop quantitative metrics for these types of risks, a food safety assessment is considered appropriate for this issue. Individual companies need to make the determination as to the level of hazard after considering the following risk factors: the concentration and volume of fecal matter, frequency of animals (observed or indicators) in the field, density of animal population and surrounding area risk – all identified during a risk assessment. A trained food safety professional should be involved in decisions related to animal intrusion. See Appendix B for more details on the qualifications for this person. Appendix B describes in detail the process used to develop these metrics 	

1223

1224

1225

TABLE 6. CROP LAND AND WATER SOURCE ADJACENT LAND USE

Land Use/Water Source	Metric (This distance may be either increased or decreased depending on risk and mitigation factors.)	Considerations for Risk Analysis*		
		Risk/Mitigation Factors	Increase Distance	Decrease Distance
Composting Operations (manure or animal products)	Due to the lack of science at this time, an interim guidance distance of 400 ft from the edge of crop is proposed. This number is subject to change as science becomes available. The proximate safe distance depends on the risk/mitigation factors listed to the right. Evaluate risk and document consideration of these factors. Research is being proposed to study appropriate distance.	Distance from active compost operation	--	--
		Topography: Uphill from crop	√	
		Topography: Downhill from crop		√
		Opportunity for water run off through or from composting operations	√	
		Opportunity for soil leaching	√	
		Presence of physical barriers such as windbreaks, diversion ditches, vegetative strips		√
Concentrated Animal Feeding Operations (as defined in 40 CFR 122.23)	Due to the lack of science at this time, an interim guidance distance of 400 ft from the edge of crop is proposed. This number is subject to change as science becomes available. The proximate safe distance depends on the risk/mitigation factors listed to the right. Evaluate risk and document consideration of these factors. Research is being proposed to study appropriate distance. <u>Distance from a CAFO is not sufficient to address/manage all potential hazards that may be associated with growing leafy greens in proximity to a CAFO. Due to the lack of science at this time, an interim guidance distance of 1200 ft. from the edge of a CAFO larger than 1000 head is established. This number is only a reference and subject to change as science becomes available. The proximate safe distance depends on many risk mitigation factors. This distance may increase or decrease after assessing the risk, determining and deploying mitigation measures and consulting with customers.</u>	Fencing and other physical barriers such as berms, diversion ditches and vegetated strips can be employed to prevent intrusion of domestic animals, control runoff, etc.		√
		Topography: Uphill from crop	√	
		Topography: Downhill from crop		√
		Opportunity for water run off through or from CAFOs	√	
		Opportunity for soil leaching		√
		Manure Management Program utilized		√

Formatted: Font: (Default) Times New Roman, 10 pt

Land Use/Water Source	Metric (This distance may be either increased or decreased depending on risk and mitigation factors.)	Considerations for Risk Analysis*		
		Risk/Mitigation Factors	Increase Distance	Decrease Distance
Non-synthetic Soil Amendment Pile (containing manure or animal products)	<p>Due to the lack of science at this time, an interim guidance distance of 400 ft from the edge of crop is proposed. This number is subject to change as science becomes available.</p> <p>The proximate safe distance depends on the risk/mitigation factors listed to the right. Evaluate risk and document consideration of these factors. Research is being proposed to study appropriate distance.</p> <p>For non-synthetic crop treatments that have been heat treated using a validated process an interim guidance distance of 30 feet from the edge of the crop is proposed</p>	Access and review COA for materials in question.		✓
		Topography: Uphill from crop	✓	
		Topography: Downhill from crop		✓
		Opportunity for water run off through or from non-synthetic soil amendment storage areas	✓	
		Opportunity for soil leaching	✓	
		Covering on pile to prevent wind dispersion		✓
Grazing Lands/Domestic Animals (includes homes with hobby farms, and non commercial livestock)	30 ft from the edge of crop.	Fencing and other physical barriers such as berms, diversion ditches and vegetated strips can be employed to prevent intrusion of domestic animals, control runoff, etc.		✓
		Topography: Uphill from crop	✓	
		Topography: Downhill from crop		✓
		Opportunity for water run off through or from grazing lands	✓	
		Opportunity for soil leaching	✓	
Homes or other building with a septic leach field.	30 ft from the edge of crop to the leach field.	Active leach field: < 10 yrs old		✓
		Active leach field: > 25 yrs old	✓	
		Inactive leach field		✓
		Topography: Uphill from crop	✓	
		Topography: Downhill from crop		✓
		Physical barriers		✓
Well Head Distance from Untreated Manure	200 ft separation of untreated manure from wells, although less distance may be sufficient.	Topography: Uphill from manure		✓
		Topography: Downhill from manure	✓	

Land Use/Water Source	Metric (This distance may be either increased or decreased depending on risk and mitigation factors.)	Considerations for Risk Analysis*		
		Risk/Mitigation Factors	Increase Distance	Decrease Distance
		Opportunity for water runoff from or through untreated manure to well head	√	
		Opportunity for soil leaching	√	
		Presence of physical barriers such as windbreaks, diversion ditches, vegetative strips		√
Surface Water Distance from Untreated Manure	At least 100 feet separation for sandy soil and 200 feet separation for loamy or clay soil (slope less than 6%; increase distance to 300 feet if slope greater than 6%) is recommended.	Topography: Uphill from manure		√
		Topography: Downhill from manure	√	
		Opportunity for water runoff from or through untreated manure to surface waters.	√	
		Opportunity for soil leaching	√	
		Presence of physical barriers such as windbreaks, diversion ditches, vegetative strips		√
Rationale	<ul style="list-style-type: none"> The bases for these distances above is best professional judgment of authors, contributors, and expert reviewers to prevent potential cross-contamination from adjacent land uses, taking into consideration the 200 foot distance cited in FDA (US FDA 2001) for separation of manure from wellheads and the 30 foot turn-around distance for production equipment. Because of the numerous factors that must be taken into account to determine appropriate distances, a qualitative assessment of the relative risk from various types of land use and surface waters was used to determine appropriate distances. 			

1226 *Producers should check for local, state and federal laws and regulations that protect riparian habitat, restrict removal of vegetation or habitat, or restrict

1227 construction of wildlife deterrent fences in riparian areas or wildlife corridors. Producers may want to contact the relevant agencies (e.g., the Regional Water

1228 Quality Control Board and state and federal fish and wildlife agencies) to confirm the details of these requirements.

1229 17. **TRANSPORTATION**

1230 When transporting lettuce/leafy greens on the farm or from the farm to a cooling, packing, or processing
1231 facility, manage transportation conditions to minimize the risk of contamination. Food contact surfaces on
1232 transportation equipment and in transporter vehicle cargo areas that are not properly maintained are potential
1233 sources of contamination.

1234 **The Best Practices Are:**

1235 • Visually inspect all shipping units and equipment used to transport leafy greens on the farm or from
1236 the farm to a cooling, packing, or processing facility to ensure they are:
1237 ○ In good, working condition; and
1238 ○ Clean before use in transporting lettuce/leafy greens.

1239 18. **DETAILED BACKGROUND GUIDANCE INFORMATION**

1240

1241 **Required Reference Documents**

1242 1. FDA Guide to Minimize Microbial Food Safety Hazards for Fresh Fruits and Vegetables
(www.foodsafety.gov/~-dms/prodguid.html)
1243 2. UFFVA Food Safety Auditing Guidelines: Core Elements of Good Agricultural Practices for Fresh
1244 Fruits and Vegetables
1245 3. UFFVA Food Safety Questionnaire for Fresh Fruits and Vegetables
1246 4. National GAPs Program Cornell University: Food Safety Begins on the Farm: A Grower Self-
1247 Assessment of Food Safety Risks

1248 **References**

1249

1250 1251 CCR Title 14 - Chapter 3.1 - Article 5. 2007. *Article 5. Composting Operation and Facility Siting and*
1252 *Design Standards*. Accessed February 15, 2007.
<http://www.ciwmb.ca.gov/regulations/Title14/ch31a5.htm#articles>
1253 Fukushima H, Hoshina K, and Gomyoda M. 1999. Long-term survival of shiga toxin-producing
1254 Escherichia coli O26, O111, and O157 in bovine feces. *Applied and environmental microbiology*
1255 65 (11):5177-81.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=10543842
1256 Gagliardi JV and Karns JS. 2000. Leaching of Escherichia coli O157:H7 in diverse soils under various
1257 agricultural management practices. *Applied and environmental microbiology* 66 (3):877-83.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=10698745
1258 Islam M, Doyle MP, Phatak SC, Millner P, and Jiang X. 2004. Persistence of enterohemorrhagic
1259 Escherichia coli O157:H7 in soil and on leaf lettuce and parsley grown in fields treated with
1260 contaminated manure composts or irrigation water. *Journal of food protection* 67 (7):1365-70.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15270487
1261 Jiang X, Morgan J, and Doyle MP. 2003. Fate of Escherichia coli O157:H7 during composting of bovine
1262 manure in a laboratory-scale bioreactor. *Journal of food protection* 66 (1):25-30.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12540177
1263 Solomon EB, Pang HJ, and Matthews KR. 2003. Persistence of Escherichia coli O157:H7 on lettuce
1264 plants following spray irrigation with contaminated water. *Journal of food protection* 66

1276 (12):2198-202.
1277 http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=14672213

1279 Stine SW, Song I, Choi CY, and Gerba CP. 2005. Application of microbial risk assessment to the
1280 development of standards for enteric pathogens in water used to irrigate fresh produce. *Journal of*
1281 *food protection* 68 (5):913-8.
1282 http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15895721

1284 Suslow TV, Oria MP, Beuchat LR, Garrett EH, Parish ME, Harris LJ, Farber JN, Busta FF. 2003.
1285 Production practices as risk factors in microbial food safety of fresh and fresh-cut produce.
1286 *Comprehensive Reviews in Food Science and Food Safety* 2S:38-77.

1287 Takeuchi K and Frank JF. 2000. Penetration of Escherichia coli O157:H7 into lettuce tissues as affected
1288 by inoculum size and temperature and the effect of chlorine treatment on cell viability. *Journal of*
1289 *food protection* 63 (4):434-40.
1290 http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=10772206

1292 Takeuchi K, Matute CM, Hassan AN, and Frank JF. 2000. Comparison of the attachment of Escherichia
1293 coli O157:H7, Listeria monocytogenes, Salmonella typhimurium, and Pseudomonas fluorescens to
1294 lettuce leaves. *Journal of food protection* 63 (10):1433-7.
1295 http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11041147

1297 Takeuchi K, Hassan AN, and Frank JF. 2001. Penetration of Escherichia coli O157:H7 into lettuce as
1298 influenced by modified atmosphere and temperature. *Journal of food protection* 64 (11):1820-3.
1299 http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11726166

1301 US EPA. 1996. *Soil Screening Guidance: Technical Background Document*. EPA/540/R95/128: Office of
1302 Solid Waste and Emergency Response, United States Environmental Protection Agency.
1303 http://rais.ornl.gov/homepage/SSG_nonrad_technical.pdf

1304 US EPA. 2002. *Implementation Guidance for Ambient Water Quality Criteria for Bacteria: May 2002*
1305 *Draft*. EPA-823-B-02-003: United States Environmental Protection Agency.
1306 <http://www.epa.gov/waterscience/standards/bacteria/bacteria.pdf>

1307 US FDA. 2001. Chapter II: Production Practices as Risk Factors in Microbial Food Safety of Fresh and
1308 Fresh-Cut Produce. In *Analysis and Evaluation of Preventive Control Measures for the Control*
1309 *and Reduction/Elimination of Microbial Hazards on Fresh and Fresh-Cut Produce*; pp.
1310 <http://www.cfsan.fda.gov/~comm/ift3-2a.html>

1311 US FDA. 2004. Federal Food, Drug, and Cosmetic Act. <http://www.cfsan.fda.gov/~lrd/cfr110.html>

1312 Wachtel MR, Whitehand LC, and Mandrell RE. 2002a. Association of Escherichia coli O157:H7 with
1313 pre-harvest leaf lettuce upon exposure to contaminated irrigation water. *Journal of food protection*
1314 65 (1):18-25.
1315 http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11808792

1317 Wachtel MR, Whitehand LC, and Mandrell RE. 2002b. Prevalence of Escherichia coli associated with a
1318 cabbage crop inadvertently irrigated with partially treated sewage wastewater. *Journal of food protection*
1319 65 (3):471-5.
1320 http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=1899045